



## **Carpal Tunnel Syndrome Caused by Anatomic Anomalies Muscles: A Three Cases Report**

**Mohamed Ali Sbai<sup>1</sup>, Rami Ben Arab<sup>1\*</sup>, Leila Essid<sup>1</sup>, Amira Gallas<sup>1</sup>  
Khaled Khelil<sup>1</sup>, Monia Boussem<sup>2</sup> and Riadh Maalla<sup>3</sup>**

<sup>1</sup>*Department of Orthopedic and Trauma Surgery, Mohamed Taher Al Maamouri Hospital, Nabeul, Tunisi, Tunisia.*

<sup>2</sup>*Department of Emergency, Mongi Slim Hospital, La Marsa, Tunisia.*

<sup>3</sup>*Department of Plastic Surgery, Rabta Hospital, Tunis, Tunisia.*

### **Authors' contributions**

*This work was carried out in collaboration among all authors. Author RBA designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors MAS and LE managed the analyses of the study. Author KK managed the literature searches. All authors read and approved the final manuscript.*

### **Article Information**

#### Editor(s):

(1) Dr. Wagih Mommtaz Ghannam, Professor, Department of General Surgery, Faculty of Medicine, Mansoura University, Egypt.

#### Reviewers:

(1) Jose Luis Turabian, Health Center Santa Maria de Benquerencia, Spain.

(2) P. K. Hota, Kaloji Narayana Rao University of Health Sciences, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50669>

**Received 17 June 2019**

**Accepted 06 August 2019**

**Published 21 August 2019**

**Case Study**

### **ABSTRACT**

Carpal tunnel syndrome (CTS) is the most frequent peripheral compression neuropathy. Anatomic variations may be encountered during carpal tunnel surgery. 1–3 Compression of the median nerve at the wrist is frequently encountered. Carpal tunnel syndrome usually occurs without any obvious extrinsic causes; several cases have however been reported caused by anomalous or hypertrophic muscles. A survey of the literature shows that compression neuropathy of the median nerve has been reported in relation with anomalies affecting three muscles: the first (or second) lumbrical, the palmaris longus and its anatomic variants and the superficial flexor of long fingers. We can suspect the presence of such an anomalous muscle when the compression syndrome concerns a patient who is not within the "usual" age group with symptoms initiated or aggravated by physical exercise. This report presents three cases of carpal tunnel syndrome caused by anatomic anomalous muscles diagnosed peroperatively.

\*Corresponding author: E-mail: [benarabrami@gmail.com](mailto:benarabrami@gmail.com);

*Keywords: Median nerve; nerve compression; carpal tunnel syndrome; anomaly; muscle; hand surgery.*

## 1. INTRODUCTION

Carpal tunnel syndrome (CTS) is a frequently encountered condition in middle-aged women; it is in most cases idiopathic. In all other "atypical" carpal tunnel syndromes, a more extensive search for external causes of the compression is necessary. An extrinsic structure is usually responsible for the compression and a systematic search for compressing structures is required [1]. Compression of the median nerve by muscle anomaly is exceptional and performs a specific symptomatology: Affects a young adult, manual worker, aggravated by manual activity. Clinical examination searches paresthesia during bending of MP. The ultrasound and MRI are used to make the diagnosis. The lifting of the median nerve compression by opening the carpal tunnel can relieve the patient [2]. We present in this work 3 cases of secondary carpal tunnel syndrome caused by muscle abnormality. The aim of this work is to present three rare cases of carpal tunnel syndrome causes.

## 2. CASES

### 2.1 Case 1

A 25-year-old female who suffered from paraesthesia, and numbness in the median nerve territory of the left hand for several years. Her symptoms were aggravated by exercise. She had a positive Phalen test and Tinel's sign at the carpal tunnel without thenar atrophy. The clinical diagnosis of CTS was confirmed by electrophysiological examination. Under loco regional anaesthesia, carpal tunnel release was performed through the classic incision. After dividing the transverse carpal ligament, an aberrant tendon was discovered on the anterior surface of the median nerve within its investing tissue (Fig. 1). The tendon was inserted deeply into the palmar aponeurosis and evoke abnormal palmaris longus tendon: palmaris profundus. The median nerve was congested. The palmaris longus tendon was palpable above the wrist crease. Further exploration above the wrist was not indicated clinically and the origin of the palmaris profundus was not established. After a few weeks the patient's symptoms resolved completely.

### 2.2 Case 2

A 35-year-old left-handed woman, without significant pathological history suffering from numbness, tingling, pain, and weakness in the left hand and affecting the thumb, the index and the middle finger, developed 6 months ago without any traumatic or microtraumatic antecedent [3]. Interrogation of the patient was recorded a progressive evolution over several months with gradual acroparaesthesia in the median nerve territory associated with palmar pain. Her symptoms were aggravated by exercise. Physical examination discovered a decrease in epicritic sensitivity with the Weber test, without evidence for motricity impairment, using preserved prehensile strength and digital winding as readouts. No abnormalities were seen on laboratory studies.


Functional exploration with electromyography identified a sensitive injury downstream of the carpal tunnel. Standard X-rays did not find abnormalities. Surgery was performed under loco-regional anaesthesia with an axillary block. Carpal tunnel release was performed through the classic incision. Abnormal lumbrical tendon was easily exposed after skin incision and opening of the mid-palmer fascia and the flexor retinaculum. The median nerve was flattened (Fig. 2) Postoperatively, the patient went to home with scar care each other day and a prophylactic treatment for algoneurodystrophy with vitamin C for one month. No immobilization was used for this purpose. Complete healing occurred after 21 days of treatment. The last clinical review at the two-month postoperative, showed an entire disappearance of the acroparaesthesia and total functional recovery including full mobility of the fingers, absence of pain, and restoration of the grip strength. No specific rehabilitation was necessary.

### 2.3 Case 3


A 65-year-old female who has suffered from paraesthesia, and numbness in the median nerve territory of the right hand for five years. Her symptoms were aggravated by exercise. Tinel's sign and Phalen test were positive. There was no evidence of thenar atrophy. The clinical diagnosis of CTS was confirmed by electrophysiological examination. Under loco regional anaesthesia, carpal tunnel release

was performed through the classic incision. After dividing the transverse carpal ligament, we identified a muscle in the carpal tunnel. Pull on the muscle led to proximal interphalangeal


joint flexion of middle finger (Fig. 3). The median nerve was congested. After a few weeks the patient's symptoms resolved completely.



**Fig. 1. Peroperating view showing palmaris profundus tendon with compression of median nerve**



**Fig. 2. Peroperating view showing hypertrophy of lumbrical tendon and compression of median nerve**



**Fig. 3. Peroperating view showing hypertrophic and engaged the flexor digitorum superficialis of the middle finger into the carpal canal**

### 3. DISCUSSION

Carpal tunnel syndrome caused by a space occupying lesion is rare and more complicated than idiopathic carpal tunnel syndrome. Compression of the median nerve with anatomic anomalies muscles is very rare and produces an atypical clinical presentation. Facing anatomic anomalies during carpal tunnel surgery is not uncommon. In a study of 382 patients, Tountas et al. reported that 38 hands had an anomalous anatomy. Still and Kleinert [4] reported a series of 9 cases of which 8 presented with a neuro-compression: A carpal tunnel syndrome in 6 cases, the anomalies involved the palmaris longus in 4 cases, the flexor digitorum superficialis in two and abnormal lumbricals in another two cases.

The pathogenesis of carpal tunnel syndrome is varied. In most cases no specific cause can be disclosed; these cases are usually termed "idiopathic" or "primary". Secondary carpal tunnel syndrome has been caused by various conditions. Abnormal or aberrant muscles have been well described, particularly in manual workers. These anomalous muscles might be hypertrophic or abnormal lumbricals, a hypertrophic flexor digitorum superficialis or an abnormal palmaris longus (profundus or reversed). All these muscles compromise the available space in the carpal canal, resulting in compression of the median nerve [5].

#### 3.1 Lumbrical Muscle

The first or second lumbrical muscle is usually held responsible for the compression.

The relationship of the lumbrical muscles and the carpal tunnel was investigated by Cobb et al. [6]. They found on cadavers that incursion of the lumbricals into the carpal canal during flexion of the fingers is a normal occurrence. It may be a possible cause of occupation-related carpal tunnel syndrome [7,8].

#### 3.2 M. flexor Digitorum Superficialis

Anomalies of the muscle belly of the flexor digitorum superficialis of long fingers as a cause for compression of the median nerve were reported and well [9,10]. In two of them the muscle belly was hypertrophic and engaged into the carpal canal, compressing the median nerve [11]; in the others an abnormal muscle belly originated in the palm and inserted onto the

superficial flexor tendon. Resection of this muscle belly seems to be necessary in such cases.

#### 3.3 Abnormal Palmaris Longus Tendon

The palmaris longus muscle is the most variable muscle in the forearm. It can be absent (15% of all patients, more on the left side and in women), hypertrophic, reversed (muscle belly distal rather than proximal), centrally placed, digastric, duplicated or bifid. It can be located deep to the transverse ligament (palmaris profundus) [12,13]. Anomalies in origin and insertion have also been described and particular accessory insertion slips are well recognized (palmaris accessorius). Lindly and Kleinert [4] found (5.7%) anomalies in 526 CTS surgeries and there was one palmaris profundus anomaly in their experience. Palmaris profundus was first described in 1908. Reimann et al. [14] found one example in 530 cadaver arms. It arises as a separate muscle in the middle third of the forearm, deep to the superficialis muscles. Its tendon courses deep into the carpal tunnel as a tenth tendon that may stray from a dorsal to palmar position within the canal, inserting into the palmar aponeurosis. The palmaris profundus is not a variation of the palmaris longus. Several reports indicate that the palmaris longus coexists with palmaris profundus [5,15].

Ultra-sonogram, CT and MRI has been performed. Recently, arthroscope or minimal invasive surgery has been preferred, however, in cases such as this involving space occupying lesions, symptoms do not improve unless open transverse carpal ligament release is performed in conjunction with removal of the SOL. Based on this study, it is important to perform special tests such as CT or MRI to identify the SOL through physical examination when unilateral CTS patients exhibit swelling or tenderness of volar wrist crease area. The use of MRI scans is an excellent way to examine soft tissue, and also has the advantage to make it possible diagnose exact location and border of lesion and also involvement of surrounding tissue. However, CT scans are superior to MRI with regard to detection of bony lesion. Ultra-sonogram has its advantages such as wide availability, lower cost, and shorter examination time; however, accurate characteristics of SOL cannot be identified [16].

However, imaging is not routinely done for carpal tunnel syndrome, and surgeon's first encounter likely will be intraoperatively. The literature is not conclusive as to whether the anomalous muscle

can cause clinical nerve compression. In most cases, the authors have advocated resection of the anomalous muscle [17]. There are reports where the patients are managed with just release of the carpal ligament, without resection of the muscle [18], or even with physical therapy alone. However, it must also be noted that, the resection of these anomalous muscles did not lead to any residual deficits. Some authors suggest excision of the muscle when there is a suspicion of it causing the compression, only after identifying the existence of normal structures to preserve function [19]. In our cases, conventional open transvers carpal ligament without resection of these anomalous muscle gave immediate painless.

#### 4. CONCLUSION

All space-occupying lesions can compromise the free course of a nerve in an anatomically limited tunnel, for anomalous, aberrant or hypertrophied muscles. A thorough knowledge of every possible anatomic variation is indispensable to surgeons who perform CTS surgery, as an appreciation of the possibilities increases the safety of the procedure [20]. In patients, not belonging to the so-called typical population for carpal tunnel syndrome, younger patients with a neurocompression syndrome related to physical activity, the treating physician should be aware of the possibility of such pathology. An anatomic anomaly may be the aetiology of the CTS and failure to address it results in the persistence to the symptoms and ends in failure to the treatment.

#### CONSENT

As per international standard or university standard, patient's consent has been collected and preserved by the authors.

#### ETHICAL APPROVAL

It is not applicable.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Chen CH, Wu T, Sun JS, Lin WH, Chen CY. Unusual causes of carpal tunnel

2. syndrome: Space occupying lesions. *J Hand Surg Eur.* 2012;37(1):14-9.
2. Sbai MA, Benzarti S, Msek H, Boussen M, Khorbi A. Carpal tunnel syndrome caused by lipoma: A case report. *The Pan African Medical Journal.* 2015;22:51.
3. Afshar, A. Palmaris profundus as the cause of carpal tunnel syndrome. *Archives of Iranian Medicine.* 2009;12(4):415-416.
4. Lindley SG, Kleinert JM. Prevalence of anatomic variations encountered in elective carpal tunnel release. *J Hand Surg Am.* 2003;28(5):849-55.
5. De Smet L. Median and ulnar nerve compression at the wrist caused by anomalous muscles. *Acta Orthop Belg.* 2002;68(5):431-8.
6. Cobb TK, An KN, Cooney WP, Berger RA. Lumbrical muscle incursion into the carpal tunnel during finger flexion. *J Hand Surg Br.* 1994;19(4):434-8.
7. Carroll M, Montero C. Rare anomalous muscle cause of carpal tunnel syndrome. *Orthop. Rev.* 1980;9:83-85.
8. Butler B. Jr, Bigley E. C. Jr. Aberrant index (first) lumbrical tendinous origin associated with carpal-tunnel syndrome. A case report. *J Bone Joint Surg Am.* 1971;53(1): 1602.
9. Hutton P, Kernohan J, Birch R. An anomalous flexor digitorum superficialis indicis muscle presenting as carpal tunnel syndrome. *Hand.* 1981;13(1):85-6.
10. Neviasser RJ. Flexor digitorum superficialis indicis and carpal tunnel syndrome. *Hand.* 1974;6(2):155-6.
11. Smith RJ. Anomalous muscle belly of the flexor digitorum superficialis causing carpal-tunnel syndrome. Report of a case. *J Bone Joint Surg Am.* 1971;53(6):1215-6.
12. Brones MF, Wilgis EFS. Anatomical variations of the palmaris longus, causing carpal tunnel syndrome: Case reports. *Plastic and Reconstructive Surgery.* 1978; 62(5):798-800.
13. Tountas CP, MacDonald CJ, Bihle DM. Carpal tunnel syndrome. A review of 507 patients. *Minn Med.* 1983;66:479-482.
14. Reimann AF, Daseler ED, Anson BJ, Beaton LE. The palmaris longus muscle and tendon. A study of 1600 extremities. *Anat Rec.* 1944;89:495-505.
15. Dyreby JR, Engber WD. Palmaris profundus rare anomalous muscle. *J Hand Surg [Am].* 1982;7:513-514.
16. Kang HJ, Jung SH, Yoon HK, Hahn SB, Kim SJ. Carpal tunnel syndrome caused

- by space occupying lesions. Yonsei Med J. 2009;50(2):257–261.
17. Bakriga B, Amouzou KS, Ayouba G, Kombate, N, Dellahn Y, Walla A, et al. Anomalous muscles: A rare cause of carpal tunnel syndrome: A case report. International Journal of Orthopaedics. 2017;4(1):705-07.
  18. Javed S, Woodruff M. Carpal tunnel syndrome secondary to an accessory flexor digitorum superficialis muscle belly: Case report and review of the literature. Hand (NY). 2014;9(4):554-55.
  19. Laxminarayan Bhandari, Michelle Palazzo. Anomalous muscles encountered in the carpal tunnel: A report of two cases. Journal of Clinical and Diagnostic Research. 2017;PD03-PD04.
  20. Suk JI, Walker FO, Cartwright MS. Ultrasonography of peripheral nerves. Curr Neurol Neurosci Rep. 2013;13(2): 328.

---

© 2019 Sbai et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

*<http://www.sdiarticle3.com/review-history/50669>*