



# Investigation of Heavy Metal Contamination in Fruits and Vegetables from Local Markets of Chennai, Tamil Nadu, India

Kalpana Palani <sup>a</sup>, Akila Srinivasan <sup>a</sup>, M. Shankar <sup>b</sup>,  
Sinduja Murugaiyan <sup>a\*</sup>, Sathya Velusamy <sup>a</sup>,  
S. Kalimuthupandian <sup>a</sup> and M. R. Ramasubramaniyan <sup>a</sup>

<sup>a</sup> National Agro Foundation, Taramani, Chennai, Tamil Nadu, India.

<sup>b</sup> Indian Agricultural Research Institute, ICAR, New Delhi, India.

## Authors' contributions

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i91287>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121554>

**Original Research Article**

**Received: 08/06/2024**

**Accepted: 14/08/2024**

**Published: 22/08/2024**

## ABSTRACT

Certain areas of Chennai such as red hills, Minjur, T. Nagar, Thiruvanmyur, Ambattur have been found to be particularly susceptible to high pollution levels due to severe traffic congestions, the ignition of coal, cement industries, State Industries Promotion Corporation of Tamil Nadu Limited (SIPCOT) and other chemical industries etc., which pose threat to the environment. The population besides these industries areas also increasing due to the employment opportunities and other activities. Considering the persistent nature, cumulative behaviour, and probability toxicity of the

\*Corresponding author: E-mail: [seethasinduja@gmail.com](mailto:seethasinduja@gmail.com);

**Cite as:** Palani, Kalpana, Akila Srinivasan, M. Shankar, Sinduja Murugaiyan, Sathya Velusamy, S. Kalimuthupandian, and M. R. Ramasubramaniyan. 2024. "Investigation of Heavy Metal Contamination in Fruits and Vegetables from Local Markets of Chennai, Tamil Nadu, India". *Journal of Advances in Biology & Biotechnology* 27 (9):179-91. <https://doi.org/10.9734/jabb/2024/v27i91287>.

heavy metals from these industries, it can enter the food chain linking the plant system to the soil (fruits & vegetables) and transformed to human body. The heavy metals, Chromium, Cadmium, Nickel and Lead, when exceed their permissible limit ( $\leq 0.05$  ppm), it causes metabolic malfunctions to human beings. At this at the background, a study was attempted on the levels of metal deposits from industries in the fruits and vegetables from the surrounding local markets in the designated locations. Although there have been numerous research studies conducted worldwide on the subject, this is the first of its kind study in Chennai district. Vegetable and fruit samples collected from the local markets near the industrial zones and in congested areas were analysed and compared for the levels against the global safety standards to ensure that fruits and vegetables can be consumed without the risk of heavy metal contamination. The study's findings demonstrated that all of the fruits and vegetables from the local markets in the study had levels of contaminants below detectable thresholds (0.05 ppm) and within WHO allowed limits (Cd- 0.2ppm, Pb - 0.3 ppm, Ni - 67.9 ppm, Cr- 0.00 ppm). This indicates there may be no possible health concerns linked to the concentrations of heavy metals in fruits and vegetables from the markets near the highly polluted industrial zones or in the areas with severe traffic congestion.




Image 1. Graphical abstract

**Keywords:** Industrial areas; heavy metals; fruits and vegetables; Chennai market.

## 1. INTRODUCTION

Population increases, urbanisation, and industrial activity lead to contamination and a detrimental influence on land, water and atmosphere [1]. Due to their long-term toxicity effects, heavy metals are considered the soil pollutants originating via natural and artificial sources. They are the primary focus of recent investigations due to the contaminating effect [2]. World over, the safety of food is a key societal concern. The increased demand for food safety over the past few decades has prompted research on the hazards related to consuming foods contaminated by

toxic substances. In regions with high industrial activity, metal concentration in soils tends to increase. These places have a few times more metal deposition more than (0.05 ppm) than uncontaminated sites [3]. The fact that pollutants may move via soils to plant to reach the food chain. Usage or direct intake by animals that consume them is the most significant effect of pollution on environmental health [4].

Traditional channels and anthropogenic mechanisms are the both ways that heavy metals enter the environment. They can be associated with the causes, such as mining

operations, deterioration of the earth's crust, and soil deterioration, liquid waste, sewage from cities, and insecticides [5]. Long-term use of pesticide use, untreated municipal effluents usage in agriculture fields, sludge waste with metal contamination are the main contributing factors to the accumulation of heavy metal in productive soils [6]. Their movement from soil to plant parts is significantly impacted by a nature of soil and plant characteristics that control their bioavailability. These substances have the potential to contaminate the atmosphere, streams, and soils from both point and non-point sources (Fig. 2) [7]. Vegetable and fruits contamination with heavy metal draws increasing attention as understanding the problem in relation to population [8]. Vegetables and fruits accumulate heavy metals as a result of air and soil contamination. The interaction with contaminated soils or air pollution can cause natural food contamination [9]. Fruits and vegetables constitute 90% of the human diet are the most common way that people are exposed to this contamination the remaining 10% is ingested through skin contact and contaminated dust inhalation [10].

The heavy metals, Cu, Ni, Fe, Ni, Cd, Hg and Co, are necessary for humans and play significant role in several metabolic processes. These metals can be harmful to both people and other living things [11]. Even at low quantities, non-essential substances including lead, cadmium, mercury, and arsenic is harmful for humans. Vegetables can accumulate heavy metals due to land and atmosphere pollution [12]. Bioaccumulation in food causes serious health risks to people. The human body absorbs these metals through food and breathing. Soil movement and its availability contribute to this heavy metal contamination [13]. Every person should eat 300 g of greens every day, including tubers, roots, and green leafy vegetables,

according to Indian Council of Medical Research Expert Committee [14]. People consume a variety of vegetables each day, depending on the produce's growing region, the type of irrigation source utilised, and its consumer age [15]. In humans, in all forms of deposition of heavy metals (HMs) results in a variety of difficulties, including renal failure, peripheral neuropathy, lung cancer, skin cancer, osteoporosis, bone thinning, and prostate cancer [16]. Heavy metal soil additions have detrimental consequences on food, crop growth, and environmental safety [17].

In Tamil Nadu, cash crops, vegetable crops and fruits crops are cultivated, 10% of all fruits and vegetables are produced in India are grown in Tamil Nadu; these include bananas, mangoes, pineapple, guava, citrus fruits, grapes, and other vegetables [18]. No investigation has found evidence of heavy metal build up in fruits or vegetables; instead, previous studies have mostly concentrated on a limited number of specific vegetable varieties [19] and on the exposure to contaminant from industrial regions such as smelters, oil refineries, petrochemical refineries, pesticide plants, cement plants, coal burning industries throughout local available markets throughout the world (Table 1) but no significant investigation is being conducted in the Chennai district. So, the main objective of the research is aimed at the heavy metal accumulation (Lead, Cadmium, Chromium and Nickel) in the fruits and vegetables from the local markets near the industrial sites and localities with severe traffic congestion namely red hills, Minjur, T. Nagar, Thiruvannamur, Ambattur in Chennai district. The findings of these trace elements were compared with standards and also investigated the everyday intake and risks of metal exposure to human health, so that vegetables and fruits can be consumed without any harm to human body.

**Table 1. Observations on heavy metals in fruits and vegetables**

S.No.	Study area	Fruit & vegetable	Ni	Cd	Cr	Pb	References
1.	Netherlands	Apple, apricot, lemon, orange	-	-	-	0.020 mg kg <sup>-1</sup> (dry weight)	[20]
2.	India	Apple, Grapes, Orange	-	-	-	224.4 ug l <sup>-1</sup> (juice)	[20]
3.	Maharashtra	mango	-	-	-	-	[20]
4.	Uttar Pradesh	Tomato	0.89	0.14	-	0.46	[16]
5.	Vellore	onion	-	0.073	-	-	[21]
6.	Nigeria	-	8.90	3.08	-	8.90	-

## 2. MATERIALS AND METHODS

### 2.1 Location of the Research

The examination was conducted in the Chennai district of Tamil Nadu, in five different parts of the city: Red Hills (13.186 and the longitude is: 80.199), Minjur, (13.269, and the longitude is 80.264), T. Nagar (13.0405026 and the longitude is: 80.2336924), Thiruvanmyur (12.982684 Latitude and 80.263369 longitude), and Ambattur (Latitude: 13.092, Longitude: 80.166). Leading industries in the region include State Industries Promotion Corporation of Tamil Nadu Limited (SIPCOT), the cement, coal, chemical, automotive, and polymer industries, among others. The city has a climate that is both wet and dry. Herewith maps (Fig. 1) displaying, sampling location relative positions (Table 2). Along with the dense population, industrial production activities have been operating in this area since the late 1970s that influence the surrounding environment. The air pollution levels

are also high with reference to PM 2.5 [22]. While being transported from the production site to the market, these fruits and vegetables are covered with polyethylene sheets; nevertheless, when they are being marketed, they are exposed to the urban environment pollutant too.

### 2.2 Sample Collection

From the areas of Thiruvanmyur, T. Nagar, Ambattur, Red hills, Minjur of Chennai districts, same type of vegetable and fruit specimens have been purchased through several nearby marketplaces (Table 2). Four samples were purchased from each market and combined at each sampling location. The collected sample (1 district x 5 local markets x 11 samples) (total n = 55 samples) were tagged, placed in Poly propylene (PP) containers, and shipped to the testing facility. The samples were properly cleaned with deionized water then dried for one day at 60°C in an oven, glass mortar was used to




Fig. 1. Study area


Table 2. Features of the Chennai market sample sites

S.No.	Study sites	Type of markets	Land uses
1.	Thiruvanmyur	Local	HT
2.	T. Nagar	Local	HT
3.	Ambattur	Local	NIA
4.	Red Hills	Local	NIA
5.	Minjur	Local	NIA

HT- Heavy traffic; NIA- Near Industrial Area

**Table 3. Botanical description of the vegetable and fruits used in the study**

S.No.	Local name	Botanical description
1.	Grapefruit	<i>Vitis vinifera</i>
2.	Apple	<i>Malus pumila</i>
3.	Guava	<i>Psidium guajava</i>
4.	Orange	<i>Citrus sinensis</i> ,
5.	Amla	<i>Phyllanthus emblica</i>
6.	Lemon	<i>Citrus limon</i>
7.	Beans	<i>Phaseolus vulgaris</i>
8.	Tomato	<i>Solanum lycopersicum</i>
9.	Potato	<i>Solanum tuberosum</i>
10.	Carrot	<i>Daucus carota</i>
11.	Cucumber	<i>Cucumis sativus</i>



**Fig. 2. Sources of heavy metals**

ground the dried samples into a homogeneous mixture, and a 2 mm nylon screen was used to filter out any coarse particles. The specimens with botanical characterizations are in Table 3.

### 2.3 Analysis of Fruits and Vegetable

For the analysis of vegetables and fruits, a 100 ml acid-washed beaker containing one gram of material was filled with 15 ml of a di-acid mixture. It is digested at 80 °C until a transparent solution was obtained. After cooling, the digested samples were filtered through Whatman No. 42 filter paper, and the filtrate was subsequently diluted to make 50 ml using deionized water in a volumetric flask. Using AAS (Atomic Absorption Spectroscopy), the heavy metals were found in the vegetable filtrate and air deposition. Using the standard solutions of varied concentrations of the appropriate heavy metals, the instrument was calibrated. Air served as the support and

acetylene gas served as the fuel. Each instance involved the work of an oxidising flame, which reveals various settings for the instrument depending on the metal.

### 2.4 Method Validation

The analytical process was validated using various metrics, such as accuracy, precision, and limit of detection.

### 2.5 Daily Intake of Heavy Metals through Fruits and Vegetables

The daily intake of heavy metals through the consumption according to the equation has been calculated as per given below formula:

$$\text{Daily intake of heavy metals } \mu\text{g day} = [\text{Daily fruit or vegetable consumption} \times \text{fruit or Vegetable heavy metal concentration}]$$

### 3. RESULTS AND DISCUSSION

As a result of their toxicity, heavy metals are regarded as the most significant environmental pollutant (Figs. 3 & 4). Table (4) illustrates the

number of heavy metals in a few fruits and vegetables from five different Chennai, Tamil Nadu marketplaces, along with a comparison to the WHO's permitted guideline (Table 5) for heavy metals.




Fig. 3. Heavy metal accumulation in fruits and vegetables from the environment




Fig. 4. Heavy metal absorption in plants and transportation to fruits and vegetables

**Table 4. Concentration of heavy metals in vegetables**

Vegetables & fruits	Location																			
	T. Nagar				Ambattur				Red Hills				Minjur			Tiruvanmyur				
	Ni	Cr	Cd	Pb	Ni	Cr	Cd	Pb	Ni	Cr	Cd	Pb	Ni	Cr	Cd	Pb	Ni	Cr	Cd	Pb
Beans	1.86	BDL	BDL	BDL	2.24	BDL	BDL	BDL	3.14	BDL	BDL	BDL	3.45	BDL	BDL	BDL	1.57	BDL	BDL	BDL
Tomato	13.1	BDL	BDL	BDL	16.2	BDL	BDL	BDL	16.32	BDL	BDL	BDL	15.22	BDL	BDL	BDL	10.3	BDL	BDL	BDL
Potato	18.3	BDL	BDL	BDL	21.5	BDL	BDL	BDL	20.2	BDL	BDL	BDL	26.14	BDL	BDL	BDL	11.6	BDL	BDL	BDL
Carrot	14.2	BDL	BDL	BDL	16.3	BDL	BDL	BDL	17.25	BDL	BDL	BDL	1.12	BDL	BDL	BDL	1.3	BDL	BDL	BDL
Cucumber	2.61	BDL	BDL	BDL	2.19	BDL	BDL	BDL	2.05	BDL	BDL	BDL	1.47	BDL	BDL	BDL	1.49	BDL	BDL	BDL
Grapes	1.87	BDL	BDL	BDL	2.13	BDL	BDL	BDL	1.44	BDL	BDL	BDL	3.12	BDL	BDL	BDL	1.63	BDL	BDL	BDL
Apple	1.54	BDL	BDL	BDL	1.33	BDL	BDL	BDL	1.38	BDL	BDL	BDL	1.02	BDL	BDL	BDL	2.01	BDL	BDL	BDL
Guava	1.88	BDL	BDL	BDL	1.42	BDL	BDL	BDL	1.47	BDL	BDL	BDL	1.33	BDL	BDL	BDL	1.24	BDL	BDL	BDL
Orange	2.26	BDL	BDL	BDL	3.52	BDL	BDL	BDL	0.87	BDL	BDL	BDL	3.28	BDL	BDL	BDL	BDL	BDL	BDL	BDL
Amla	1.22	BDL	BDL	BDL	1.17	BDL	BDL	BDL	0.98	BDL	BDL	BDL	1.74	BDL	BDL	BDL	BDL	BDL	BDL	BDL
Lemon	2.33	BDL	BDL	BDL	1.97	BDL	BDL	BDL	1.24	BDL	BDL	BDL	0.68	BDL	BDL	BDL	BDL	BDL	BDL	BDL

**Table 5. Limit of heavy metals permitted in fruits and vegetables**

S.No.	Element	FAO/WHO (mg/Kg)
1.	Cd	0.2
2.	Pb	0.3
3.	Ni	67.9
4.	Cr	-

\*FAO: Food and Agriculture Organization \*WHO: World Health Organization

### 3.1 Lead Concentration in Fruits and Vegetables


Lead could enter the human system through food, drink, and air. No known safe level of exposure exists. It is mostly stored in livers and kidneys. The toxicity of daily lead intakes is very low (0.05) at the determined concentration levels [23].

According to dry weight of the sample, the amounts of Pb were estimated. The Pb concentrations results with below detectable limits (BDL  $\leq 0.05$ ) for all the vegetable and fruit samples of all the sites (Table 4) which is well within the 0.3 mg/kg permissible range (Table 5) (Fig. 5). The levels of lead identified in a variety of greens collected from markets in Tanzania, Saudi Arabia, Nigeria, and Serbia were less than the highest amounts allowed (0.30 and 0.10 mg kg<sup>-1</sup> for greens) [24] and also with the vegetable field Nigerian urban wastewater that was used for irrigation contained a permissible amount of

Pb level within an acceptable threshold [25]. Air, water, and food all have pathways for lead to get into the human circulatory system. But this decreased level of lead in the study area is because, Pb is the least transportable element from the soil, and soil Pb concentrations are highly correlated with minerals, oxides of manganese, hydroxyls ions of iron and aluminium, and organic matter in soils [26]. Whereas the soils near the industrial areas could not be rich in Pb is not organic matter and hence the insufficient level of Pb [27]. The daily intake of Pb is also at BDL from the study area of the selected vegetables and fruits. Also, the lead concentration in the mango fruit was less than the instrument detection limit, indicating that either no lead was present in the sample or only a trace amount was [28]. According to numerous reports, using whole fruits-including their peels-carries a risk of consuming heavy metals that may be found in the peel [29]. The evaluation of health risks is within the hazard quotient level with less than 1 as per [30].

**Table 6. The calories of fruit contribute for heavy metal**

S.No.	Heavy metal	The calories of fruit contribute for an average person
1.	Pb	36.89 g
2.	Cd	5.54 g
3.	Zn	0.205 mg
4.	Cu	0.288 mg
5.	Co	45.24 g
6.	Ni	0.116 mg



**Fig. 5. Chromatogram showing the concentration of Pb**



### 3.2 Nickel Concentration in Fruits and Vegetables

Based on the collected samples of vegetables and fruits, nickel concentration shown in (Table 4) but all the concentrations were below the permissible limit of Ni (67.9 mg/Kg) in all the samples from the study area. This corroborated with the findings of [31] where the Ni concentration in selected fruits and vegetables in Pakistan had 3-12 mg Ni kg<sup>-1</sup> and also the Ni levels varied between 0.05 and 0.24 mg/Kg in Jews mallow and highest observed in fluted pumpkin plant 0.01 - 0.1 to 0.23 and 0.25. However, Ni level of 0.067 mg/Kg for Indian Basil which is within the range of values obtained from this study. Wastewater irrigation of vegetables in Morocco also resulted in 85 mg Ni kg<sup>-1</sup> [32]. In India, vegetables grown in peri-urban areas and irrigated with untreated wastewaters had Ni concentration as high as 506 mg Ni kg<sup>-1</sup> [33]. Ni in the soil at less than pH 6.7 is in soluble form and most Ni exists as insoluble hydroxides. So, the Ni concentration would be in insoluble form so that the uptake is less [34].

### 3.3 Cadmium Concentration in Fruits and Vegetables

Extended cadmium accumulation in humans can result in several health problems, including kidney damage, renal disorders, and human carcinogens [16]. Due to cadmium build-up in the body, prolonged ingestion of several of these fruits and vegetables causes health issues [35]. Even at low concentrations, Cadmium is an extremely hazardous metal which forms up in internal organs and could result in fatal conditions [36]. The level of Cd is determined to be BDL ( $\leq 0.05$ ) in all the vegetables and fruits from the sites taken for the study area (Table 4) which falls in the acceptable range of 0.2 mg/Kg (Table 5) which correlates with the findings of [36] that the majority of the three regions fruits and vegetables had cadmium concentrations that were within the acceptable range of 0.2 mg/kg [37] and also the vegetables and fruits spices analysed for the Cd in Nigeria were found to be within the limit. The results are in line with the research undertaken in different parts of the world (Table 1). Furthermore, this is a result of physiological barriers that prevent harmful heavy metal(loid)s from entering phloem, being loaded or unloaded, or accumulating in fruits [38] were found to be under the World Health Organisation/ Food and Agricultural Organisation (WHO/FAO) and Indian acceptable limits [8] and (Table 5).

Another important criterion for low Cd transportation into the produce is certain soil characteristics, including soil pH and organic matter that can have an impact on the sequences of Cd speciation in soils [39]. Increased pH results in decreased concentration of Cd in plants due to decreased Cd availability in soils [2]. The organic matter in alkaline industrial waste water intruded soils easily forms soluble Cd complexes and prevents exchange of ions and transfer to roots because most of the industries of study area were observed to let alkaline effluents into the soils and water nearby.

### 3.4 Chromium Concentration in Fruits and Vegetables

Cr (VI) and its metabolites, especially chromate, enter the human body through different ways (inhalation, ingestion and skin contact), which will cause pathological changes to the human organs and systems (respiratory tract, skin, gastrointestinal tract and so on) and even increase the incidence rate and mortality of many cancers [40]. Prolonged human exposure can cause gastrointestinal upset, respiratory problems, kidney and liver damage, and altered genetic material, among other conditions [41]. At modest levels, Cr (VI) can be acutely hazardous and is recognised as a known human respiratory carcinogen. The carcinogenic risk factor (CRs) value of Nickel (Ni), Chromium (Cr) and Cadmium (Cd) were found higher than the tolerance limit ( $< 10^{-4}$ ). The range of CRs for Cr from 0.0046 to 0.02104 in Bhoothnath vegetable market (BVM, 0.0067 to 0.0325 in Mohanlal Ganj vegetable markets MVM and 0.00416 to 0.040 in South city vegetable market (SVM) and The CRs value of Cr and shown the high potential of carcinogenesis through the consumption of the vegetables having high Cr level [15].

Within the selected fruits, the Cr concentrations were found to be BDL in the study area and also within the permissible limit (Table 4) Pomelo and star fruit had the highest chromium values, at 0.217 mg kg<sup>-1</sup> respectively but Mangoes from the Chattogram region and jackfruit from both regions showed no signs of chromium. Therefore, the local population is safe and no such urgent measurements are needed [42] because of even at low concentrations the Cr is toxic in nature. Exposure to Cr can cause lung, kidney, or liver damage. The uptake is BDL ( $\leq 0.05$ ) in the selected vegetables and fruits may be due to the chromium behaviour in ion

exchange capacity in soil. Since Cr is less mobile, due to its adsorption rate, Cr<sup>6+</sup> reduces as pH rises [43].

### 3.5 Daily Intake of Vegetables and Fruits

The provisional tolerated daily intake (PTDI) is typically used to express consumer exposure and the associated health hazards. The WHO has established a heavy metal intake threshold determined by the typical adult's body weight, is 60 kg. Vegetable and fruit intake per person per day is 98 and 78 grams, respectively. From the results of the study, all the samples were found to have below detectable limit and do not pose any threat to humans. When compared with the permissible limits. For an average person, the calories of fruit contribute heavy metal intake (Table 6), respectively. Therefore, heavy metal contamination in industrial area is more likely than food crops cultivated in isolated locations. Fruits and vegetables that had been treated had lower heavy metal contents [44]. Since the heavy metal pose detrimental influence on human health as well as the nutritional properties of agricultural goods; they must comply within WHO's specified safety criteria and there must be sufficient control over the levels in food.

## 4. CONCLUSION

The findings of the research revealed that concentrations of toxic heavy metals were below detectable levels of Cd, Ni, Cr and Pb in all the fruits and vegetables samples collected from the study area. According to this study, the heavy metals were not present in the produce coming from the research zone because of the considerable traffic around the market sites, vehicle emissions, smoke, and dust from nearby factories. The hazards related to toxicity of contaminants in vegetables and fruits should be taken into account for the local people, as these vegetables and fruits are among the most popular nourishment consumed in the study area. Further research may be carried out extensively and periodically in the markets near the other industrially polluted areas of Tamil Nadu to ascertain the food safety levels as per the norms laid down by the Food Safety and Standards Authority of India (FSSAI).

### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models

(ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

## ACKNOWLEDGEMENTS

The authors thank National Agro Foundation, Chennai, for providing all research resources.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Dev LK, K, NB, S, Philips M. Heavy Metals and Microplastics as a Function of Environmental Pollution: A Comprehensive Study of a Ramsar Site in Chennai, Tamilnadu India. Saravanan and NB, Srilakshmi and Philips, Mayola, Heavy Metals and Microplastics as a Function of Environmental Pollution: A Comprehensive Study of a Ramsar Site in Chennai, Tamilnadu India.
2. Wang L, Rinklebe J, Tack FM, Hou D. A review of green remediation strategies for heavy metal contaminated soil. *Soil Use and Management*. 2021;37(4):936-963.
3. Long Z, Huang Y, Zhang W, Shi Z, Yu D, Chen Y, et al. Effect of different industrial activities on soil heavy metal pollution, ecological risk, and health risk. *Environmental Monitoring and Assessment*. 2021;193:1-12.
4. Uddin MM, Zakeel MCM, Zavahir JS, Marikar FM, Jahan I. Heavy metal accumulation in rice and aquatic plants used as human food: A general review. *Toxics*. 2021;9(12):360.
5. Parida L, Patel TN. Systemic impact of heavy metals and their role in cancer development: A review. *Environmental Monitoring and Assessment*. 2023; 195(6):766.
6. Gavrilescu M. Water, soil, and plants interactions in a threatened environment. *Water*. 2021;13(19):2746.
7. Mawari G, Kumar N, Sarkar S, Daga MK, Singh MM, Joshi TK, Khan NA. Heavy metal accumulation in fruits and vegetables and human health risk assessment: Findings from Maharashtra,

- India. Environmental Health Insights. 2022;16:11786302221119151.
8. Xiang M, Li Y, Yang J, Lei K, Li Y, Li F, et al. Heavy metal contamination risk assessment and correlation analysis of heavy metal contents in soil and crops. Environmental Pollution. 2021;278:116911.
  9. Najmi A, Albratty M, Al-Rajab AJ, Alhazmi HA, Javed SA, Ahsan W, et al. Heavy metal contamination in leafy vegetables grown in Jazan region of Saudi Arabia: Assessment of possible human health hazards. International Journal of Environmental Research and Public Health. 2023;20(4):2984.
  10. Mitra S, Chakraborty AJ, Tareq AM, Emran TB, Nainu F, Khusro A, et al. Impact of heavy metals on the environment and human health: Novel therapeutic insights to counter the toxicity. Journal of King Saud University-Science. 2022;34(3): 101865.
  11. Bansal OP. Accumulation of potentially toxic metals and pesticides in fish other seafood and their impact on human. International Journal of Recent Scientific Research. 2021;12(8):42848-42868.
  12. Alengebawy A, Abdelkhalek ST, Qureshi SR, Wang MQ. Heavy metals and pesticides toxicity in agricultural soil and plants: Ecological risks and human health implications. Toxics. 2021;9(3):42.
  13. Mukherjee A, Dutta S, Goyal TM. India's Phytonutrient Report-A Snapshot of Fruits and Vegetables Consumption, Availability and Implications for Phytonutrient Intake (No. id: 12064); 2017.
  14. Kumar P, Kumar S, Singh RP. High contamination of toxic heavy metals in vegetables and their associated health risk assessment from different vegetable markets of the metropolitan city, Lucknow, India. International Journal of Environmental Research. 2021;15:837-847.
  15. Gupta N, Yadav KK, Kumar V, Krishnan S, Kumar S, Nejad D, et al. Evaluating heavy metals contamination in soil and vegetables in the region of North India: Levels, transfer and potential human health risk analysis. Environmental Toxicology and Pharmacology, 2021;82: 103563.
  16. Tefera M, Teklewold A. Health risk assessment of heavy metals in selected Ethiopian spices. Heliyon. 2021;7(5).
  17. Saravanakumar V, Malaiarasan U, Balasubramanian R, Angles S. Production efficiency and profitability of major farming systems in Tamil Nadu. Agricultural Economics Research Review. 2020;33(confspl):99-108.
  18. Ganesh KS, Sridhar A, Vishali S. Utilization of fruit and vegetable waste to produce value-added products: Conventional utilization and emerging opportunities-A review. Chemosphere. 2022;287:132221.
  19. Širić I, Eid EM, El-Morsy MH, Osman HE, Adelodun B, Abou Fayssal S, Kumar P. Health risk assessment of hazardous heavy metals in two varieties of mango fruit (*Mangifera indica* L. var. Dasher and Langra). Horticulturae. 2022;8(9):832.
  20. Bharani V, Sathyaraj S, Sinha A, Sugankumar D, Osborne JW. A survey on the accumulation of chromium in plants near various tannery industries of Vellore district, TN. Research Journal of Pharmacy and Technology. 2015;8(9): 1205-1208.
  21. Jia X, Fu T, Hu B, Shi Z, Zhou L, Zhu Y. Identification of the potential risk areas for soil heavy metal pollution based on the source-sink theory. Journal of Hazardous Materials. 2020;393:122424.
  22. Ara A, Usmani JA. Lead toxicity: A review. Interdisciplinary Toxicology. 2015;8(2): 55.
  23. Hussain S, Rengel Z, Qaswar M, Amir M, Zafar-ul-Hye M. Arsenic and heavy metal (cadmium, lead, mercury and nickel) contamination in plant-based foods. Plant and Human Health, Volume 2: Phytochemistry and Molecular Aspects. 2019;447-490.
  24. Ogunkunle ATJ, Bello OS, Ojofeitimi OS. Determination of heavy metal contamination of street-vended fruits and vegetables in Lagos state, Nigeria. International Food Research Journal. 2014;21(6).
  25. Li Q, Wang Y, Li Y, Li L, Tang M, Hu W, et al. Speciation of heavy metals in soils and their immobilization at micro-scale interfaces among diverse soil components. Science of the Total Environment. 2022; 825:153862.

26. Stefanowicz AM, Kapusta P, Zubek S, Stanek M, Woch MW. Soil organic matter prevails over heavy metal pollution and vegetation as a factor shaping soil microbial communities at historical Zn–Pb mining sites. *Chemosphere*. 2020;240:124922.
27. Pant N, Subedee A, Gharti RB, Shrestha S. Spectrometric determination of heavy metals present in mango fruit of Nepali origin. *Amrit Research Journal*. 2020;1(1):72-77.
28. Czech A, Malik A, Sosnowska B, Domaradzki P. Bioactive substances, heavy metals, and antioxidant activity in whole fruit, peel, and pulp of citrus fruits. *International Journal of Food Science*. 2021;2021:1-14.
29. Zhang A, Cortes V, Phelps B, Van Ryswyk H, Srebotnjak T. Experimental analysis of soil and mandarin orange plants treated with heavy metals found in Oilfield-Produced wastewater. *Sustainability*. 2018; 10(5):1493.
30. Bukhari IH, Ramzan M, Riaz MUHAMMAD, Bokhari TH, Rehman GHANA, Munir SHAHIDA. Determination of trace heavy metals in different varieties of vegetables and fruits available in local market of Shorkot Pakistan. *Int J Curr Pharm Res*. 2013;5(2):101-105.
31. AL-Jaboobi M, Zouahri A, Tijane M, El Housni A, Mennane Z, Yachou H, et al. Evaluation of heavy metals pollution in groundwater, soil and some vegetables irrigated with wastewater in the Skhirat region “Morocco”. *Journal of Materials and Environmental Science*. 2014;5(3):961–966.
32. Yadav A, Yadav PK, Shukla DN. Investigation of heavy metal status in soil and vegetables grown in urban area of Allahabad, Uttar Pradesh, India. *International Journal of Scientific and Research Publications*. 2013;3(9):1-7.
33. Bhalerao SA, Sharma AS, Poojari AC. Toxicity of nickel in plants. *Int. J. Pure Appl. Biosci*. 2015;3(2):345-355.
34. Rahimzadeh MR, Rahimzadeh MR, Kazemi S, Moghadamnia AA. Cadmium toxicity and treatment: An update. *Caspian Journal of Internal Medicine*. 2017;8(3):135.
35. Genchi G, Sinicropi MS, Lauria G, Carocci A, Catalano A. The effects of cadmium toxicity. *International Journal of Environmental Research and Public Health*. 2020;17(11):3782.
36. Ezeilo CA, Okonkwo SI, Chibuzor C, Onuorah ALC, Ugwunnadi NE. Determination of heavy metals in some fruits and vegetables from selected market's in Anambra state. *ACTA Sci. Nutr. Heal*. 2020;4:163-171.
37. Nazir R, Khan M, Masab M, Rehman HU, Rauf NU, Shahab S, et al. Accumulation of heavy metals (Ni, Cu, Cd, Cr, Pb, Zn, Fe) in the soil, water and plants and analysis of physico-chemical parameters of soil and water collected from Tanda Dam Kohat. *Journal of Pharmaceutical Sciences and Research*. 2015;7(3):89.
38. Li Z, Liang Y, Hu H, Shaheen SM, Zhong H, Tack FM, Zhao J. Speciation, transportation, and pathways of cadmium in soil-rice systems: A review on the environmental implications and remediation approaches for food safety. *Environment International*. 2021;156:106749.
39. Christou A, Georgiadou EC, Zissimos AM, Christoforou IC, Christofi C, Neocleous D, Fotopoulos V. Uptake of hexavalent chromium by tomato (*Solanum lycopersicum* L.) plants and mediated effects on their physiology and productivity, along with fruit quality and safety. *Environmental and Experimental Botany*. 2021;189:104564.
40. Ahmed F, Fakhrudin ANM, Fardous Z, Chowdhury MAZ, Rahman MM, Kabir MM. Accumulation and Translocation of Chromium (Cr) and Lead (Pb) in Chilli Plants (*Capsicum annum* L.) Grown on Artificially Contaminated Soil. *Nature Environment & Pollution Technology*. 2021; 20(1).
41. Islam MA, Parvin M, Quader MFB, Hossain MS. Determination of toxic heavy metal contents of some selected tropical fruits grown in industrial areas of Bangladesh. *Dhaka University Journal of Science*. 2022;70(1):22-27.
42. Bamigboye OS. Exploration for Fe-Mn oxides using geochemical signatures in soil: A Case Study of Part of Northwestern Nigeria. In *Geochemistry*. Intech Open; 2021.
43. Chen YG, Huang JH, Luo R, Ge HZ, Wołowicz A, Wawrzkiwicz M, et al. Impacts of heavy metals and medicinal

- crops on ecological systems, environmental pollution, cultivation, and production processes in China. *Ecotoxicology and Environmental Safety*. 2021;219:112336.
44. Abdurrahman MI, Chaki S, Saini G. Stubble burning: Effects on health & environment, regulations and management practices. *Environmental Advances*. 2020; 2:100011.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/121554>