



# Phase Transfer Catalysis: New Technology to Boost Organic Farming

**Narayan G. Hegde** <sup>a++\*</sup>

<sup>a</sup> Nature Cure Ashram, Urulikanchan, Pune, India.

## **Author's contribution**

*The sole author designed, analysed, interpreted and prepared the manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/ajrcs/2024/v9i3292>

**Open Peer Review History:**  
This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/122540>

**Review Article**

**Received: 03/07/2024**  
**Accepted: 05/09/2024**  
**Published: 09/09/2024**

## **ABSTRACT**

Introduction of Phase Transfer Catalysis (PTC) technology, which was developed in the 1970s has revolutionised the dyes and pharmaceutical industries in many ways. PTC is a process that facilitates the organic compounds to move from one phase to another phase, without any change in the chemical qualities. Application of this technology was explored in the 1980s for agriculture, by using different plant extracts as PT catalysts. The objective was to facilitate increased absorption of nutrients by plants, by stimulating plant growth and easy availability of nutrients in the soil. Introduction of nano-carbon materials with nutrients and herbal growth stimulants could further support the crop growth by facilitating increased uptake of nutrients, apart from enabling the plants to develop resilience to harsh environmental conditions, resulting in higher crop yields. Some organic bio-stimulants developed recently by Snehasrusthi Group in India have been very successful in improving the growth, yield and quality of several seasonal and perennial crops in India. These bio-stimulants have excellent potential to boost organic agricultural production and income of farmers in the future.

<sup>++</sup>Managing Trustee;

<sup>\*</sup>Corresponding author: E-mail: [narayanhegde47@gmail.com](mailto:narayanhegde47@gmail.com);

**Cite as:** Hegde, Narayan G. 2024. "Phase Transfer Catalysis: New Technology to Boost Organic Farming". *Asian Journal of Research in Crop Science* 9 (3):86-94. <https://doi.org/10.9734/ajrcs/2024/v9i3292>.

**Keywords:** Phase transfer catalysis; PTC; organic farming; plant growth stimulants.

## 1. INTRODUCTION

Phase Transfer Catalysis is a new chemical process invented in the recent years. In synthetic organic chemistry, the major challenge has been to blend two chemicals of different nature to form a new product without allowing any intermittent reactions to alter their qualities. A common example is to mix a water soluble reagent and another non-water insoluble organic reagent. This problem is generally solved by adding a solvent that is both water-like and organic-like, which holds both the chemicals together. The efficiency of such blending is high when the catalyst has a larger area of interface. This method of promoting reactions between reagents with opposite solubility preferences is known as Phase Transfer Catalysis (PTC) technology and the chemical used to facilitate such reaction is known as Phase Transfer Catalyst. This mechanism was first conceived in the 1950s by Henry Hennis, a process chemist at Dow Chemical [1]. He outlined the extraction mechanism to explain a successful alkylation reaction which he developed and that was catalysed by a quaternary ammonium salt [2]. 10 years later, Charles Starks, who was associated with Continental Oil, reinvented the extraction mechanism and provided necessary support with detailed kinetic studies. Starks coined the term "Phase-Transfer Catalysis", and published a paper in the Journal of the American Chemical Society in 1971 [3]. Since then, PTC has become a practical methodology for organic synthesis. Now PTC is a prime synthetic tool, widely used for industrial applications and manufacturing process. This technology is now being applied effectively in a wide range of organic reactions, in the synthesis of pharmaceuticals, agricultural chemicals, flavouring agents, perfumes, dyes and for polymerization reactions, monomer

synthesis and polymer modifications, for environmental control processes and for many other applications [4].

The objective of this paper is to review the role of PTC in different sectors of the chemical industry and how this technology could contribute to the agricultural production, particularly to reduce the use of expensive and harmful agro-chemicals and to promote residue-free, food production. The methodology adopted was to review the basic principle involved in making use of PTC technology by different industries and how some efforts were made by different scientists to apply this technology for promoting plant growth, without affecting food safety and quality. Therefore the study included the review of research papers and industrial processes on PTC and personal interactions with the scientists and chemical professionals engaged in industrial production of organic chemicals. Field visits were also carried out to verify the response of various agricultural crops to newly generated organic compounds in producing residue-free, organic agricultural produce.

## 2. TYPES OF CATALYSTS USED FOR PTC

The principle of PT catalysis is based on the ability of certain phase-transfer catalysts to facilitate the transport of one reagent from one phase into another un-mixable phase wherein the other reagent exists. Thus, reaction is made possible by bringing together the reagents which are originally in different phases. However, it is necessary that the transferred species is in an active state for effective PT catalytic action, and that it is regenerated during the organic reaction [5].

**Table 1. Commonly used PT catalysts**

| Catalysts         | Cost                          | Stability and Activity   | Use and Recovery   |
|-------------------|-------------------------------|--|--|
| Ammonium salts    | Cheap                         | Moderately stable in basic conditions and up to 100°C. Decomposition by Hofmann elimination under basic conditions. Moderately active. | Widely used. Recovery is relatively difficult.   |
| Phosphonium salts | Expensive than ammonium salts | Thermally more stable than ammonium salts, although less stable under basic conditions.  | Widely used. Recovery is relatively difficult.   |
| Crown ethers      | Expensive                     | Stable and very active catalysts both under basic conditions and at higher temperatures up to even 150-200°C.                          | Often used. Recovery is difficult and causes environmental issues due to their toxicity. |

| Catalysts | Cost       | Stability and Activity  | Use and Recovery   |
|-----------|------------|---|--|
| Cryptands | Expensive  | Stable and highly reactive, excluding the presence of strong acids. | Used sometimes despite high costs and toxicity, due to higher reactivity.            |
| PEG       | Very cheap | Lower activity but more stable than quaternary ammonium salts.      | Often used; Larger quantities of catalyst cause no problems. Fairly easy to recover. |

Source: Jagtapa and others [2]

There are many types of PTCs, such as quaternary ammonium and phosphonium salts, crown ethers, cryptands, etc. The types of catalysts used and their details are presented in Table 1.

Among these, the quaternary ammonium salts are the cheapest and hence most widely used in the industry [6-11]. Despite the development of PTC in organic synthesis, the mechanistic aspects of PTC remain unclear, mainly due to the difficulty of investigating biphasic systems and many complex parameters which are required to be analysed.

### 3. MECHANISM OF EXCHANGE

The mechanism of PTC reaction was first proposed by Dr. Charles Starks, who called this mechanism as the Extraction Mechanism [12]. According to him, a quaternary ammonium halide (Q+X-) present in the aqueous phase undergoes anion exchange with the anion (Y-) of the reactant dissolved in the aqueous solution. The ion pair formed (Q+Y-) can pass through the liquid-liquid interface as a result of its lipophilic nature, and goes from the interface into organic phase. The cycle continues with subsequent return of catalyst to the aqueous phase [13-16]. In the organic phase, the anion of the ion-pair is nucleophilic, which undergoes a nucleophilic substitution reaction with the organic reagent, forming the desired product. The catalyst subsequently returns to the original aqueous phase and the cycle continues. Thus, PTC is an important and efficient technique for various organic transformations due to its operational simplicity, mild reaction conditions, use of safe and inexpensive reagents and solvents, and possibility of conducting reactions on a large scale [17]. With this technology, it is feasible to facilitate the interphase migration of substrates making reactions between the reagents in two immiscible phases [18]. PTC is also a useful method for carbohydrate synthesis, as the reactions can be performed on a large scale [19].

**Choice of PT Catalyst:** There are two important requirements for using any molecule as a PT catalyst [15]. It should be cationic and must

contain an adequate chemical structure, which is capable for splitting the nucleophilic anion in the organic phase. Further, the cation-anion bonding must be low for superior anionic reactivity. Some important factors in employing a PT catalyst are stability beneath reaction conditions, ease of formation or availability of catalyst, ease of separation or recovery and non-toxicity [16-21]. Although no definite guidelines are provided for the selection of an ideal catalyst for a given reaction system, depending on some of the above factors, appropriate PT catalysts can be used. The application of PTC technology became cheaper and easily available with the use of alternative raw materials like potassium carbonate and aqueous NaOH solution, thereby avoiding the requirement of severe anhydrous conditions and expensive solvents and organometallic reagents. As PT catalyst, quaternary ammonium salts based on cinchona alkaloids are often used for asymmetric epoxidation of electron-deficient alkenes. Chiral crown ethers and guanidinium salts are also used for asymmetric epoxidation of electron-deficient alkenes [22-24]. Presently, temperature-controlled phase-transfer catalyst and hydrogen bond phase-transfer catalyst are regarded as better alternatives to traditional phase-transfer catalysts such as onium salt, crown ether and polyether phase-transfer catalyst. This is because of their excellent properties such as nontoxicity, well-cycling performance and low cost [25]. Use of this technology was further extended in the synthesis process for producing fine chemicals used as agrochemicals, pharmaceuticals, perfumery and fragrances, dyes, paper, etc. during the recent years [26-28].

### 4. APPLICATION OF PTC FOR ORGANIC FARMING

In the mid-1980s, when Indian farmers started realising the disadvantages of using higher doses of agro-chemicals, causing the pollution of soil, water and food products, apart from high cost of production, there was a strong wave of organic farming, as an alternative to high input agriculture. Most of the farmers were directionless, while the scientists were looking for

viable alternatives. The challenge was to reduce the quantity of chemical fertilizers being applied, without starving the crop for nutrients and depriving protection against pests and diseases, without using chemicals which cause environmental pollution and health hazards.

With regard to nutrients, it was a known fact that the quantity of nutrients applied in the form of chemical fertilizers was significantly higher than the actual quantity of nutrients absorbed by the crop plants. Higher doses of fertilizers affected soil pH and soil texture, reduced soil permeability and increased the salt content, thereby depleting the soil productivity. Application of fertilizers in lower doses did not maximise the crop yields, because of their inability to absorb the required quantities of nutrients. It was primarily because, a significant part of the nutrients available in the soil and fertilizers were fixed and insoluble for absorption by the plants. Secondly, the plants did not have robust vigour to absorb more nutrients. Hence the scientists were searching for two types of solutions, firstly to increase the release of nutrients and secondly to increase the absorption capacity of plants. The other challenge was to replace the chemical nutrients with organic nutrients. Any breakthrough in these areas could revolutionise the agricultural industry in many ways. With regard to plant protection, the challenge was to avoid harmful chemicals with safer botanical and safe organic products [29].

## 5. AN INITIATIVE TO TAP NEW TECHNOLOGIES FOR IMPROVING AGRICULTURAL PRODUCTION

During the same period in the 1980s, Mr. Dhanjay Dhekane, a senior organic chemist, working on pharmaceuticals and dyes in Ahmedabad, India was using PTC technology for developing new formulations. After thoroughly reading a classic book on PTC, titled 'Phase transfer catalysis: Principles and Technique' authored by Starks and Liotta, published in 1978, he was conversant about the advantages of its application in different fields. He has been using triethyl benzyl ammonium chloride, triethyl ammonium chloride and crown ethers as PT Catalysts for producing bulk drug intermediates like n-butyl di-ethyl malonate. The use of PTC technology reduced the process time from 25-30 hours to only six minutes with improvement in the quality and yield. Impressed with this technology, he was trying to apply PTC technology in other fields.

This was the period when India had taken benefits of Green Revolution for over 25 years and the crops were not responding proportionately to increased doses of chemical fertilizers. Increasing cost of inputs and incidences of pests and diseases were bothering the policy makers and farmers. Thus promotion of organic farming was a hot topic for discussion in the country. Hailing from the agriculturalist family, Mr. Dhekane was also trying to understand the problems of the farmers and explore suitable solutions. He was aware that several botanical extracts were used for boosting plant growth stimulation and pest control by farmers. He found out that the alkaloids and some other phytochemicals present in these plant extracts had certain molecular structures similar to the chemicals used as PT catalysts. So, he thought of mixing these extracts with micro-nutrients, organic nutrients like cow urine, humic acid, sea weed extract, etc. for spraying on plants of different vegetable crops. These plants responded to the treatment and exhibited higher growth. The results were outstanding. This motivated him to take up series of research trials by mixing herbal extracts of neem and drum-stick leaves, pongamia seed extract, etc. with humic acid, fulvic acid, protein hydrolysates, etc. for spraying on different crops at different stages. The results were spectacular (Personal communications). There was a significant improvement in plant growth, flowering intensity, size of the produce, increase in yield and improvement in the quality. The incidences of pests and diseases were also low. So he was convinced about the application of PTC technology for boosting agricultural production.

**New Product Development:** Mr. Dhekane went a step forward to explore the scope for helping the farmers interested in organic farming with a new and unique product. The aim was to increase the crop yield through the following approaches:

1. Increase the appetite of the plants to consume or absorb more nutrients from the soil;
2. Make more nutrients freely available for the plants to be absorbed from the soil;
3. Sustained release of nutrients to ensure steady supply, without any wastage in the soil through leaching;
4. Develop resistance or provide protection against pests and diseases;
5. Improve the yield and quality of the produce.

Focusing on the above objectives, various ingredients were introduced for different cocktails to be used for spray and soil application and tested on different seasonal and perennial crops.

**Application of Nano-carbon:** Nanocarbon was another important component, which was included in these products, because of its benefits to plant growth. It was fairly well known that nanocarbon has a significant role in increasing crop productivity through soil Improvement, slow release of nutrients and pesticides, growth stimulation, plant protection, environmental sensing, etc. Nanocarbon materials were cost-effective, and less toxic to plants, unlike metal and polymers. They were also helpful in maintaining the equilibrium in the environment by targeted and sustained release of fertilizers and pesticides [30]. Nanocarbon materials were immensely beneficial as biosensors, carriers, and light convertors, which could improve the sustainability of agriculture. Various benefits of nano-carbon for agriculture are presented in Table 2 [31].

**Support for Green Harvest:** Based on the field performance, Mr. Dhekane successfully developed four new generation bio-stimulants for improving agricultural production, for the first time, by incorporating certain ingredients to

facilitate the phase transfer catalysis and sustained release catalysis, which were widely used in pharmaceuticals industry. With a commitment to promote eco-friendly farming, he shifted his focus from pharmaceuticals to agriculture and established a new firm under the banner of 'Snahasrusthi' to take up commercial production of these bio-stimulants. The details of these products are presented in Table 3 [32]. These organic products were tested on a large scale through series of field trials on different crops.

The effects of these organic bio-stimulants on different crops are summarised in Table 4 (Personal communications and Field records). The results highlighted phenomenal increase in healthy vegetative growth, yield and quality, without proportional increase in the cost of inputs. These bio-stimulants have received safety approval from the concerned government authorities for application on all types of crops, which are now marketed in Maharashtra and Gujarat states and the response from farmers has been overwhelming [33]. It is anticipated that these products will immensely benefits the farmers across the country. With such new generation safe and eco-friendly inputs, bright day are ahead for the farmers and consumers.

**Table 2. Applications of nanocarbon in agriculture**

| Application area           | Nanocarbon application                 | Benefits and impacts                            |
|----------------------------|--|---|
| Soil improvement           | Nanocarbon soil additives              | Improved soil structure and nutrient retention  |
|                            | Nanocarbon-based soil amendments       | Enhanced microbial activity and fertility       |
| Nutrient management        | Nanocarbon-coated fertilizers          | Controlled nutrient release and efficiency      |
|                            | Nanocarbon nutrient carriers           | Better nutrient uptake and reduced leaching     |
| Pest control               | Nanocarbon-based pesticides            | Targeted pest control and reduced pollution     |
|                            | Nanocarbon-enhanced biopesticides      | Reduced environmental impact                    |
| Water purification         | Nanocarbon-based water filters         | Contaminant removal and improved water quality  |
|                            | Nanocarbon for heavy metal adsorption  | Cleaning polluted water sources                 |
| Crop monitoring            | Nanocarbon-enabled sensors             | Real-time data for optimized cultivation        |
|                            | Nanocarbon-enhanced imaging systems    | Detailed insights into crop health and growth   |
| Environmental remediation  | Nanocarbon-based soil remediation      | Contaminant adsorption and soil restoration     |
|                            | Nanocarbon for air and water clean-up  | Removal of pollutants and toxins                |
| Smart delivery systems     | Nanocarbon-controlled release systems  | Targeted and controlled delivery of substances  |
|                            | Nanocarbon-responsive sensors          | Environment-triggered payload release           |
| Precision agriculture      | Nanocarbon-enabled precision farming   | Tailored practices for optimized yields         |
|                            | Nanocarbon-based data-driven decisions | Enhanced resource efficiency                    |
| Customized plant nutrition | Nanocarbon nutrient release systems    | Customized nutrient delivery for plants         |
|                            | Nanocarbon-modified soil amendments    | Improved nutrient availability and plant growth |

Source: Bhojija, [31]

**Table 3. Composition of Bio-stimulants developed by 'Snehasrusthi' Group**

| Products       | Composition  | Benefits  |
|----------------|--|---|
| Soil Guard     | Activated nano-carbon (N300), Humic acid, <i>Pongamia pinnata</i> , <i>Aloe indica</i> , <i>Moringa oleifera</i> , Trace elements, PTC, Cow urine              | Increases- soil organic carbon, tolerance to salinity and climate resilience, seed germination, crop root system, nutrient uptake, crop growth and yield                |
| Bio Shield     | Activated nano-carbon (N300), Fulvic acid, <i>Pongamia pinnata</i> , <i>Aloe indica</i> , <i>Moringa oleifera</i> , Trace elements, PTC, Cow urine, and others | Improvements - in cell division. photosynthesis, plant growth, flowering, fruit set, yield and quality, shelf life, climate resilience, tolerance to pests and diseases |
| Sneha Sampurna | Activated nano-carbon (N300), PTC, <i>Sargassum wightii</i> , <i>Saccharum officinarum</i> , DSW extract   | Increase in soil fertility, plant growth, flowering, fruit set, yield and quality   |
| Sneha Shakti   | Plant extracts rich in amino acids, organic nitrogen, natural phosphorous, sulphur, zinc and other trace elements  | Improves – soil structure, soil organic carbon, water holding capacity, root system, crop yield   |

Source: Snehasrusthi, [32]

**Table 4. Performance of different crops in response to 'Snehasrusthi' Bio-stimulants**

| Agri-produce   | Bio-stimulants Used   | Results without Using 'Snehasrusthi' Products  | Results on Using 'Snehasrusthi' Products   |
|--|---|--|--|
| Banana   | Soil Guard<br>Bio Shield<br>Sneha Sampurna<br>Sneha Shakti  | Normal Bunch of Bananas, weighing about 27-30 kg. Ave. Banana size 7-9 inch.                                   | Bunch of Bananas weighing 42-45 kg. Average banana size 10-11 inch.  |
| Mango (Kesar variety in Junagadh, Gujrat)                | Soil Guard<br>Bio Shield<br>Sneha Sampurna<br>Sneha Shakti  | Each box of 10 kg contained 38-42 fruits. Size of Mangoes was decreasing from lower to upper branches on tree. | Each box of 10 kg contained 28-30 fruits. Size of Mangoes was uniform throughout the tree. Yield 60-65% higher.    |
| Mango-100 year old Alphonso trees in Deoghad Maharashtra | Soil Guard<br>Bio Shield,<br>Sneha Sampurna<br>Sneha Shakti | Trees yielded only 17-20 boxes of 5 dozen each. Shelf life: 7-9 days.  | Trees yielded about 47-50 boxes of 5 dozen each. Shelf life: 15-17 days.   |
| Lemon in Karveer block, Kolhapur, Maha                   | Soil Guard<br>Bio Shield                                    | Plants were 2.5 feet tall with smaller leaves. Yielded no fruits.  | Plant were 4-4.5 feet tall with bigger leaves. Each plant had several fruits                                       |
| Turmeric Selam variety                                   | Soil Guard<br>Bio Shield                                    | Plants were small. Yield 500-650 gm/rhizome (17-21 qt./ acre dry turmeric). Curcumin content was 3.5-3.7%.     | Bigger plants, larger leaves, yielded 2.5-4.0 kg /rhizome (dry turmeric 38-42 qt/acre). Curcumin content 4.3-4.7%. |
| Onion  | Soil Guard<br>Bio Shield<br>Sneha Sampurna<br>Sneha Shakti  | Production: 120 qt./acre. Onion size was medium  | Production: 180-200 qt./ ac. Onion were medium and large size, fresh looking                                       |
| Potato Pink variety for making chips                     | Soil Guard<br>Bio Shield<br>Sneha Sampurna<br>Sneha Shakti  | Production: 5-7 medium size potatoes/plant.  | Production: 12-14 medium size potatoes/plant.  |
| Wheat (Khapali / Emmer)                                  | Soil Guard<br>Bio Shield<br>Sneha Sampurna<br>Sneha Shakti  | Production: 12-15 quintals/ acre   | Better vegetative growth. Production: 24-25 quintals/ acre   |
| Cotton   | Soil Guard<br>Bio Shield<br>Sneha Sampurna<br>Sneha Shakti  | Yield: 8.5-8.7 qt./acre. Lesser staple length and tensile strength. Susceptible to pests.                      | Yield: 20-22 qt./acre. Better staple length and tensile strength.  |
| Sugarcane  | Soil Guard<br>Bio Shield<br>Sneha Sampurna<br>Sneha Shakti  | 7-8 shoots/plant. Cane weight 1.5-1.8 kg and length 11-12 feet. Yield: 45 tons/ac. Sugar content: 8.5-9%.      | 17-18 shoots/plant. Cane weight 2.5 kg, length 17-19 feet. Yield: 75 tons/acre. Sugar content: 9.5-9.7%.           |

Source: Unpublished Reports and personal observations

## 6. CONCLUSION

Application of Phase Transfer Catalysis (PTC) technology, which had revolutionised the dyes and pharmaceutical industries in many ways was introduced in Indian agriculture, during the mid-1980s. Herbal based bio-stimulants along with phase transfer catalysts, nano-carbon and micro-nutrients indicated their potential to boost plant growth, yield and quality. The phase transfer catalysts facilitated the increase in absorption of nutrients by stimulating plant growth and availability of nutrients in the soil. Introduction of nano-carbon particles with nutrients and herbal growth stimulants could further support agricultural crops by facilitating the increase in uptake of nutrients, apart from enabling the plants to develop resilience to harsh environmental conditions, resulting in higher crop yields. Some organic bio-stimulants developed recently in India have excellent potential to boost green agricultural production and income of farmers in the future.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during writing or editing of this manuscript.

## COMPETING INTERESTS

Author has declared that no competing interests exist.

## REFERENCES

1. Freedman HH. Industrial applications of phase transfer catalysis (PTC): Past, present and future. *Pure & Appl. Chem.* 1986;58(6):857-868.
2. Jagtapa Jayashri R, Jadhava Pranali A, Deodhara Meenakshi N, Chavana Rajshree S. Phase transfer catalysis: A green approach inorganic synthesis. *Journal for Basic Sciences.* 2022;22(11): 292-310.
3. Starks, Charles M. Phase-transfer catalysis. I. Heterogeneous reactions involving anion transfer by quaternary ammonium and phosphonium salts. *J. Am. Chem. Soc.* 1971;93(1): 195–199.

Available:<https://doi.org/10.1021/ja00730a033>

4. Ahluwalia VK, Kidwai M. Green Catalysts. In: *New Trends in Green Chemistry.* Dordrecht: Springer Netherlands. 2004;27–38.
5. Hashimoto Takuya, Maruoka Keiji. The basic principle of phase-transfer catalysis and some mechanistic aspects. In *Asymmetric Phase Transfer Catalysis.* Edited by Keiji Maruoka. WILEY-VCH GmbH & Co. KGaA, Weinheim; 2008.
6. Makosza M, Fedoryński M. Phase transfer catalysis. *Catalysis Reviews.* 2003;45(3-4):321-367.  
Available:<https://doi.org/10.1081/CR-120025537>
7. Kwak SY, Wong MH, Lew TTS, Bisker G, Lee MA, Kaplan A, et al. Nanosensor technology applied to living plant systems. *Annu. Rev. Anal. Chem.* 2017;10:113–140.
8. Giraldo JP, Wu H, Newkirk GM, Kruss S. Nano-biotechnology approaches for engineering smart plant sensors. *Nat. Nanotechnol.* 2019;14:541–553.
9. Starks CM, Liotta CL, Halpern M. Phase-transfer catalysis: Fundamentals, applications, and industrial perspectives. New York: Chapman & Hall. 1994;668.
10. Starks CM, Halpern M. Phase-transfer catalysis: Fundamentals, applications, and industrial perspectives. Springer Science & Business Media; 2012 Dec 6.
11. Anastas PT, Kirchhoff MM, Williamson TC. Catalysis as a foundational pillar of green chemistry. *Applied Catalysis A: General.* 2001 Nov 30;221(1-2):3-13.
12. Chen B, Dingerdissen U, Krauter JG, Rotgerink HL, Möbus K, Ostgard DJ, et al. New developments in hydrogenation catalysis particularly in synthesis of fine and intermediate chemicals. *Applied Catalysis A: General.* 2005 Feb 25;280(1):17-46.
13. Giraldo JP, Landry MP, Kwak SY, Jain RM, Wong MH, Iverson NM, Ben-Naim M, Strano MS. A ratiometric sensor using single chirality near infrared fluorescent carbon nanotubes: Application to in vivo monitoring. *Small.* 2015;11:3973–3984.
14. Wu H, Nibler R, Morris V, Herrmann N, Hu P, Jeon SJ, Kruss S, Giraldo JP. Monitoring plant health with near-infrared fluorescent H<sub>2</sub>O<sub>2</sub> nanosensors. *Nano Lett.* 2020;20:2432–2442.

15. Whitesides GM, Wong CH. Enzymes as catalysts in synthetic organic chemistry [new synthetic methods (53)]. *Angewandte Chemie International Edition in English*. 1985 Aug;24(8):617-38.
16. Xiao Y, Wu C, Cui P, Zhou L, Yin Q. Pursuing Green and Efficient Agriculture from Molecular Assembly: A Review of Solid-State Forms on Agrochemicals. *Journal of Agricultural and Food Chemistry*. 2023 Jul 7;71(28):10500-24.
17. Mishra M, Török B. Diastereo- and enantioselective synthesis by non-traditional activation methods: Ultrasonic, microwave, electro-, photo- and mechano-chemically activated reactions. In *Non-traditional Activation Methods in Green and Sustainable Applications*. Elsevier Inc; 2021.  
Available:<https://doi.org/10.1016/C2018-0-04332-7>
18. Xie G, Vločskó B, Török B. Green synthetic methods in drug discovery and development. In *Contemporary Chemical Approaches for Green and Sustainable Drugs Advances in Green and Sustainable Chemistry*. 2022;Chapter 8: 201-279.
19. Cormier EP, Molander GA. Synthesis: Carbon with two heteroatoms each attached by a single bond. In *Comprehensive Organic Functional Group Transformations II*. Eds. Katritzky AR, Taylor RJK. Elsevier Science; 2005.
20. Varughese P. Phase Transfer Catalysts - Principles and Techniques (Starks, Charles M.; Liotta, Charles). *J Chem Educ*. 1979;56(7):A242.
21. Wu H, Li Z. Recent advances in nano-enabled agriculture for improving plant performance. *The Crop Journal*. 2022; 10(2022):1–12.
22. Jia Z, Luo S. Asymmetric oxidative  $\alpha$ -functionalization of carbonyls Via C-O/C-N Bond Formation. In *Comprehensive Chirality (2nd Edn.)*; 2024.  
Available:<https://doi.org/10.1016/B978-0-32-390644-9.00025-1>.
23. Liu S, Zhu W. A minireview of phase-transfer catalysis and recent trends. *Biomedical Journal of Scientific & Technical Research*. 2022;45(4):36691-36702.
24. Nobuta T, Hamada S, Ueda Y, Kawabata T. Oxidation: Epoxidation of C=C. *Comprehensive Chirality*. 2024;(2nd Edn): 32-77.  
Available:<https://doi.org/10.1016/B978-0-32-390644-9.00057-3>.
25. Chandel M, Kaur K, Sahu BK, Sharma S, Panneerselvam R, Shanmugam V. Promise of nano-carbon to the next generation sustainable agriculture. *Carbon*. 2022;188(March 2022):461-481.
26. Zhu L, Chen L, Gu J, Ma H, Wu H. Carbon-Based Nanomaterials for Sustainable Agriculture: Their Application as Light Converters, Nanosensors, and Delivery Tools. *Plants (Basel)*. 2022;11(4): 511.  
DOI: 10.3390/plants11040511
27. Busacca CA, Fandrick DR, Song JJ, Senanayake CH. The growing impact of catalysis in the pharmaceutical industry. *Advanced Synthesis & Catalysis*. 2011 Aug;353(11-12):1825-64.
28. Ooi T, Kameda M, Maruoka K. Design of N-spiro C 2-symmetric chiral quaternary ammonium bromides as novel chiral phase-transfer catalysts: Synthesis and application to practical asymmetric synthesis of  $\alpha$ -amino acids. *Journal of the American Chemical Society*. 2003 Apr 30;125(17):5139-51.
29. Stepha G E J. Impact of green revolution in India. *International Journal of Health Sciences*. 2022;6(S4):5291–5297.  
Available:<https://doi.org/10.53730/ijhs.v6nS4.10077>.
30. Zahmakıran M, Özkar S. Metal nanoparticles in liquid phase catalysis; from recent advances to future goals. *Nanoscale*. 2011;3(9):3462-81.
31. Bhojiya, Ali Asger. The potential contribution of nanocarbon to fostering sustainable agriculture for future generations Nano Carbons. 2023;1(1): 228.  
Available:[https://www.researchgate.net/publication/375915828\\_The\\_potential\\_contribution\\_of\\_nanocarbon\\_to\\_fostering\\_sustainable\\_agriculture\\_for\\_future\\_generations](https://www.researchgate.net/publication/375915828_The_potential_contribution_of_nanocarbon_to_fostering_sustainable_agriculture_for_future_generations)
32. Snehasrusti Pvt. Ltd. Ahmedabad, Gujarat. Product Catalogue. 2024;26.



33. Hegde, Narayan G. Nanotechnology: and Horticultural Research. 2023;10(1):  
Cutting-edge tool for increasing agricultural 20-28.  
production. Asian Journal of Agricultural

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/122540>