



# Hepatoprotective Investigation of *Ficus lyrata* and *Ficus elastic* Leaves Extract

**Mrinal Aryan <sup>a++\*</sup>, Ashish Manigauha <sup>a#</sup> and Vishal Gupta <sup>a#</sup>**

<sup>a</sup> Faculty of Pharmacy, Mansarovar Global University Bhopal (M.P.), India.

## **Authors' contributions**

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## **Article Information**

DOI: <https://doi.org/10.9734/jpri/2024/v36i117606>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124456>

**Original Research Article**

**Received: 02/08/2024**

**Accepted: 03/10/2024**

**Published: 21/10/2024**

## **ABSTRACT**

In hepatotoxic Albino rats caused by isoniazid and paracetamol, the hydroalcoholic extract of *Ficus lyrata* and *Ficus elastic* leaves was evaluated for hepatoprotective activity. The degree of protection was measured by estimating biochemical parameters such as serum glutamate Oxaloacetate transaminase (SGOT), serum glutamate pyruvate transaminase (SGPT), ALP and the level of total serum bilirubin. In rats that had been toxicated with isoniazid and paracetamol. Hydroalcoholic extract (100 mg/kg, 200 mg/kg) demonstrated a noteworthy hepatoprotective effect in a dose-dependent manner. The extract's hepatoprotective benefits were on par with those of the prescribed medication silymarin 2.5 mg/kg body weight.

<sup>++</sup>Research Scholar;

<sup>#</sup>Professor;

<sup>\*</sup>Corresponding author: E-mail: [mrinalaryan@gmail.com](mailto:mrinalaryan@gmail.com);

**Cite as:** Aryan, Mrinal, Ashish Manigauha, and Vishal Gupta. 2024. "Hepatoprotective Investigation of *Ficus Lyrata* and *Ficus Elastic* Leaves Extract". *Journal of Pharmaceutical Research International* 36 (11):130-38. <https://doi.org/10.9734/jpri/2024/v36i117606>.

**Keywords:** *Ficus lyrata*; *Ficus elastic*; Hepatoprotective activity.

## 1. INTRODUCTION

Herbal remedies derived from nature are carefully considered to be safe and effective natural treatments for a range of illnesses. The purpose of this study was to demonstrate the hepatoprotective, antioxidant, and preliminary chemical screening properties of *Ficus elastic* and *lyrata*. After thoroughly washing the dried Leaves material under running water, an electronic grinder was used to grind it. The powder was extracted in stages using the maceration process and an ethanol-water solvent combination. *Ficus lyrata* and *Ficus elastic* were evaluated using various standards criteria, including phytochemical screening, percentage loss, yield, and organoleptic evaluation.

The India rubber plant, or *Ficus elastica*, is a huge tree in the Moraceae family that was formerly a significant supplier of subpar natural rubber. In the early 20th century, the unrelated rubber tree primarily took its position as a source of rubber. Although it is native to Southeast Asia, the India rubber plant is widely grown indoors in pots elsewhere. *Ficus lyrata* is a species of plant in the mulberry and fig family Moraceae. It is sometimes referred to as the fiddle-leaf fig, banjo fig, fiddle-leaved fig tree, lyre leaf fig tree, or lyre-leaved fig tree.

*Ficus lyrata* and *Ficus elastic* leaf are significant medicinal plants that are used to cure a variety of illnesses in traditional medicine. For two reasons, the liver may be regarded as the most significant organ in drug toxicity: first, it serves as a functional barrier between the site of absorption and the systemic circulation and is a key location for the metabolism and removal of foreign substances; second, these characteristics also make the liver a prime target for drug toxicity. Consequently, drug-induced liver damage presents a significant clinical challenge. Since the liver is essential for the detoxification and excretion of several endogenous and exogenous substances, damage to it or impairment of its function can have a number of negative effects on health. Management of liver diseases is still a challenge to modern medicine [1].

## 2. METHODOLOGY

### 2.1 Plant material

#### 2.1.1 Selection & collection of plant material [2-4]

Ethnobotanical surveys were carried out in tribal localities of Samastipur District north Bihar region.

Fresh Leaves of *Ficus lyrata* and *Ficus elastic* were collected from area adjoining forests of Samastipur District north Bihar region in the month of November.

#### 2.1.2 Drying & storage collection of plant material [5-7]

Leaves of *Ficus lyrata* and *Ficus elastic* were carried out in sun but under the shade. Leaves of *Ficus lyrata* and *Ficus elastic* were preserved in plastic bags and closed tightly and powdered as per the requirements.

### 2.2 Extraction Procedure [8,9]

100 grams of dried *Ficus lyrata* plant material and 100 grams of *Ficus elastic* were thoroughly extracted using the maceration process. Using a solvent mixture of ethanol-water (70:30), using maceration process for 48 hrs, filtered and dried using vacuum evaporator at 40°C. 70% ethanol was used to extract plant material constantly and for 48 hours at room temperature. It was proved that the extraction technique left no trace upon evaporation by adding a few drops of extract from the thimble. (Pradhan et al., 2010, Ansari 2001, Mukherjee 2007).

The following formula was used to calculate the chosen plant materials' percentage yield. The percentage yield of each extract was calculated by using following formula:

$$\text{Percentage Yield} = \frac{\text{Weight of Extract}}{\text{Weight of Powder drug taken}} \times 100$$

### 2.3 Selection of Animals

Swiss albino rats weighing between 150 and 200 g were employed in the current study. Before the trial started, the animals were given two weeks to acclimate. They were then kept in typical laboratory settings, with a temperature of 25±2°C, a relative humidity of 45–65%, and a 12-hour light/dark cycle. Throughout the investigation, the animals were given unlimited water and regular laboratory animal nutrition. The Institution Animal Ethical Committee formally accepted the protocol for animal experimentation. The Organization for Economic Co-operation and Development (OECD) guideline No. 420 was used while testing for acute oral toxicity. (CPCSEA NO.1582/PO/Re/S/11/CPCSEA) In this investigation, water was available to the Swiss

albino rats at all times during an overnight fast. For the first 24 hours and then for the following 14 days, the animals were monitored for death or any aberrant behavior after the extract was given orally at a concentration of 2000 mg/kg body weight. There were further behavioral, neurological, and autonomic reactions seen.

## **2.4 Isoniazid and Paracetamol Induced Hepatoprotective Activity of Leaves Extract of *Ficus lyrata* and *Ficus elastic*, [9-12]**

### **2.4.1 Experimental designs for *Ficus lyrata* leaves extract Isoniazid induced hepatoprotective**

Group –I: Normal control (Sterile distilled water ml/kg, p.o.)

Group –II: Isoniazid solutions were prepared in sterile distilled water (100 mg/kg, p.o.)

Group -III: *Ficus lyrata* Extract (100mg/kg, p.o.) + Isoniazid (100 mg/kg, p.o.)

Group –IV: *Ficus lyrata* Extract (200mg/kg, p.o.) + Isoniazid (100 mg/kg, p.o.)

Group –V: Silymarin (2.5 mg/kg, p.o.) + Isoniazid (100 mg/kg, p.o.)

Five groups, each with six creatures, were created from the animals. Group –I received 1ml/kg p.o. of sterile distilled water was given to the first group. Group II was received Isoniazid solutions at a dose of 100 mg/kg p.o. Group-III and Group -IV were received 100 mg/kg and 200 mg/kg of hydroalcoholic leaves extract of *Ficus lyrata*, respectively. For 21 days, Group 5 was given silymarin (2.5 mg/kg p.o.) once daily. The animal was anesthetized with ether after 21 days in order to draw blood from the retroorbital plexus, and it was subsequently slaughtered while still under ether anesthesia in order to remove the liver. Numerous biochemical analyses were conducted. (Jiang *et al.*, 2004; Saleem *et al.*, 2008).

### **2.4.2 Experimental Designs for *Ficus lyrata* leaves extract paracetamol induced hepatoprotective**

Group –I: Normal control (Sterile distilled water ml/kg, p.o.)

Group –II: Paracetamol solutions were prepared in sterile distilled water (50 mg/kg, p.o.)

Group -III: *Ficus lyrata* Extract (100mg/kg, p.o.) + Paracetamol, (50 mg/kg, p.o.)

Group –IV: *Ficus lyrata* Extract (200mg/kg, p.o.) + Paracetamol, (50 mg/kg, p.o.)

Group –V: Silymarin (2.5 mg/kg, p.o.) + Paracetamol, (50 mg/kg, p.o.)

Five groups, each with six creatures, were created from the animals. Group –I received 1ml/kg p.o. of sterile distilled water was given to the first group. Group II was received paracetamol solutions at a dose of 50 mg/kg p.o. Group-III and Group -IV were received 100 mg/kg and 200 mg/kg of hydroalcoholic leaves extract of *Ficus lyrata*, respectively. For 21 days, Group 5 was given silymarin (2.5 mg/kg p.o.) once daily. The animal was anesthetized with ether after 21 days in order to draw blood from the retroorbital plexus, and it was subsequently slaughtered while still under ether anesthesia in order to remove the liver. Numerous biochemical analyses were conducted.

### **2.4.3 Experimental designs for *Ficus elastic* leaves extract isoniazid induced hepatoprotective**

Group –I: Normal control (Sterile distilled water ml/kg, p.o.)

Group –II: Isoniazid solutions were prepared in sterile distilled water (100 mg/kg, p.o.)

Group -III: *Ficus elastic* Extract (100mg/kg, p.o.) + Isoniazid (100 mg/kg, p.o.)

Group –IV: *Ficus elastic* Extract (200mg/kg, p.o.) + Isoniazid (100 mg/kg, p.o.)

Group –V: Silymarin (2.5 mg/kg, p.o.) + Isoniazid (100 mg/kg, p.o.)

Five groups, each with six creatures, were created from the animals. Group –I received 1ml/kg p.o. of sterile distilled water was given to the first group. Group II was received Isoniazid solutions at a dose of 100 mg/kg p.o. Group-III and Group -IV were received 100 mg/kg and 200 mg/kg of hydroalcoholic leaves extract of *Ficus elastic*, respectively. For 21 days, Group 5 was given silymarin (2.5 mg/kg p.o.) once daily. The animal was anesthetized with ether after 21 days in order to draw blood from the retroorbital plexus, and it was subsequently slaughtered while still under ether anesthesia in order to remove the liver. Numerous biochemical analyses were conducted.

### **2.4.4 Experimental designs for *Ficus elastic* leaves extract paracetamol induced hepatoprotective**

Group –I: Normal control (Sterile distilled water ml/kg, p.o.)

Group –II: Paracetamol solutions were prepared in sterile distilled water (50 mg/kg, p.o.)

Group -III: *Ficus elastic* Extract (100mg/kg, p.o.) + Paracetamol, (50 mg/kg, p.o.)

Group -IV: *Ficus elastic* Extract (200mg/kg, p.o.) + Paracetamol, (50 mg/kg, p.o.)

Group -V: Silymarin (2.5 mg/kg, p.o.) + Paracetamol, (50 mg/kg, p.o.)

Five groups, each with six creatures, were created from the animals. Group -I received 1ml/kg p.o. of sterile distilled water was given to the first group. Group II was received paracetamol solutions at a dose of 50 mg/kg p.o. Group-III and Group -IV were received 100 mg/kg and 200 mg/kg of hydroalcoholic leaves extract of *Ficus elastic*, respectively. For 21 days, Group 5 was given silymarin (2.5 mg/kg p.o.) once daily. The animal was anesthetized with ether after 21 days in order to draw blood from the retroorbital plexus, and it was subsequently slaughtered while still under ether anesthesia in order to remove the liver. Numerous biochemical analyses were conducted.

### 3. RESULTS AND DISCUSSION

#### 3.1 Result of Percentage Yield

The percentage yield of leaf extracts from *Ficus lyrata* was displayed in Table 1. The ethanol-water extract had a greater yield of 12.39%, respectively. The ethanol-water leaf extract yielded 8.15%, which was comparable to the *ficus elastic* of the leaves extract.

**Table 1. Percentage yield of leaves of *Ficus lyrata* and *Ficus elastic***

S.N.	Plant Name	Percentage yield (%)
1	<i>Ficus lyrata</i>	12.39
2	<i>Ficus elastic</i>	8.15

#### 3.2 In -vivo Hepatoprotective Activity of *Ficus lyrata* & *Ficus elastic* Leaves Extract

Dose-dependent hepatoprotective effect was demonstrated by both test groups, i.e., high dose and low dosage treated groups. Liver activity improved in the test groups that received only the plant extract. It is evident that "*Ficus lyrata* and *Ficus elastic* leaves" are plants with hepatoprotective properties. According to this study, the leaves of *Ficus lyrata* and *Ficus elastica* significantly reduce the hepatotoxicity caused by the medications used to treat tuberculosis. *Ficus lyrata* and *Ficus elastic* leaves may have a hepatoprotective effect

because of their antioxidant capability, which implies that plant extract may be helpful in preventing liver damage brought on by oxidative stress.

Elevated blood levels of SGPT, SGOT, and ALP activities indicate hepatocyte injury to the liver and, in turn, indirect impairment of liver function after APAP-induced hepatotoxicity. Following APAP treatment, there was a significant ( $p < 0.05$ ) increase in SGPT, SGOT, and ALP activities, as shown in Table. *Ficus lyrata* and *Ficus elastic* leaf extract treatments at 100 and 200 mg/kg both markedly decreased the increase of these enzymes ( $p < 0.05$ ). The observed decrease in hepatic enzymes was comparable to both the control group's and the group that received silymarin pretreatment. The apparent leakage of cellular enzymes into plasma is one of the classic indicators of hepatic injury or damage.

Furthermore, the presence or lack of particular enzymes in the bloodstream can be used to determine the kind and extent of liver damage. Alkaline phosphatase, aspartate aminotransferase, and alanine aminotransferase are frequently utilized as marker enzymes in general alanine aminotransferase measurements to assess APAP-induced hepatotoxicity. This study's improvement in SGPT, SGOT, ALP, and serum bilirubin levels demonstrates the hepatoprotective impact of *Ficus lyrata* and *Ficus elastic* leaves. Serum bilirubin increases caused by isoniazid, paracetamol, and *Ficus lyrata* and *Ficus elastic* leaf extract are suppressed. Transaminase levels have been shown to rise following Isoniazid therapy in earlier investigations [13]. The increase is time-dependent, with a significant elevation observed after 48 hours ( $p < 0.05$ ), indicating severe hepatocellular damage from these typically cytoplasmic enzymes leaking into the bloodstream. (Asha, et al., 2004).

#### 3.3 Hepatoprotective Activity of *Ficus lyrata* Leaves Extract in Isoniazid Induced Hepatotoxicity

Hepatoprotective activity of hydroalcoholic extract of *Ficus lyrata* leaves and Silymarin on % SGOT levels, %SGPT levels, % serum bilirubin levels, % ALP levels in Isoniazid induced hepatotoxicity in rats Tables 1,2,3,4. Results of Hepatoprotective activity of hydroalcoholic extract of *Ficus elastic* leaves and Silymarin on % SGOT levels, %SGPT levels, % serum bilirubin levels, % ALP levels in Isoniazid induced hepatotoxicity in rats Tables 5,6,7,8.

**Table 2. Effect of Hydroalcoholic extract of *Ficus lyrata* leaves and Silymarin on % SGOT levels in Isoniazid induced hepatotoxicity in rats. [14]**

Treatment	Dose	SGOT (%)
Normal	1 ml/kg, p.o.	162 ± 2.5
Isoniazid	100 mg/kg, p.o.	341.45 ± 6.5
<i>Ficus lyrata</i> leaves	100 mg/kg p.o.	214.0 ± 3.0***
<i>Ficus lyrata</i> leaves	200 mg/kg p.o.	189.0 ± 3.5***
Silymarin	2.5 mg/kg p.o.	168.0 ± 2.9***

**Table 3. Effect of Hydroalcoholic extract of *Ficus lyrata* leaves and Silymarin on %SGPT levels in Isoniazid induced hepatotoxicity in rats [14]**

Treatment	Dose	SGPT (%)
Normal	1 ml/kg, p.o.	160.0 ± 2.0
Isoniazid	100 mg/kg, p.o.	321.0 ± 3.50
<i>Ficus lyrata</i> leaves	100 mg/kg p.o.	194.0 ± 4.0***
<i>Ficus lyrata</i> leaves	200 mg/kg p.o.	179.0 ± 3.50***
Silymarin	2.5 mg/kg p.o.	155.0 ± 3.50***

**Table 4. Effect of *Ficus lyrata* leaves and Silymarin on % serum bilirubin levels in Isoniazid induced hepatotoxicity in rats**

Treatment	Dose	Serum Bilirubin (%)
Normal	1 ml/kg, p.o.	120.0 ± 5.50
Isoniazid	100 mg/kg, p.o.	285.0 ± 3.50
<i>Ficus lyrata</i> leaves	100 mg/kg p.o.	159.0 ± 4.51***
<i>Ficus lyrata</i> leaves	200 mg/kg p.o.	129.0 ± 1.60***
Silymarin	2.5 mg/kg p.o.	115.0 ± 4.50***

**Table 5. Effect of *Ficus lyrata* leaves and Silymarin on % ALP levels in Isoniazid induced hepatotoxicity in rats**

Treatment	Dose	ALP (%)
Normal	1 ml/kg, p.o.	162.0 ± 3.20
Isoniazid	100 mg/kg, p.o.	325.0 ± 5.57
<i>Ficus lyrata</i> leaves	100 mg/kg p.o.	219.0 ± 4.30***
<i>Ficus lyrata</i> leaves	200 mg/kg p.o.	186.0 ± 3.78***
Silymarin	2.5 mg/kg p.o.	170.0 ± 5.40***

**Table 6. Effect of Hydroalcoholic extract of *Ficus elastic* leaves and Silymarin on % SGOT levels in Isoniazid induced hepatotoxicity in rats**

Treatment	Dose	SGOT (%)
Normal	1 ml/kg, p.o.	160 ± 2.5
Isoniazid	100 mg/kg, p.o.	328.45 ± 6.5
<i>Ficus elastic</i> leaves	100 mg/kg p.o.	205.0 ± 3.5***
<i>Ficus elastic</i> leaves	200 mg/kg p.o.	177.0 ± 3.5***
Silymarin	2.5 mg/kg p.o.	165.0 ± 2.5***

**Table 7. Effect of Hydroalcoholic extract of *Ficus elastic* leaves and Silymarin on %SGPT levels in Isoniazid induced hepatotoxicity in rats**

Treatment	Dose	SGPT (%)
Normal	1 ml/kg, p.o.	155.0 ± 2.0
Isoniazid	100 mg/kg, p.o.	321.0 ± 3.50

Treatment	Dose	SGPT (%)
<i>Ficus elasticleaves</i>	100 mg/kg p.o.	188.0 ± 4.0***
<i>Ficus elasticleaves</i>	200 mg/kg p.o.	165.0 ± 3.50***
Silymarin	2.5 mg/kg p.o.	162.0 ± 3.50***

**Table 8. Effect of *Ficus elasticleaves* and Silymarin on % serum bilirubin levels in Isoniazid induced hepatotoxicity in rats**

Treatment	Dose	Serum Bilirubin (%)
Normal	1 ml/kg, p.o.	120.0 ± 5.50
Isoniazid	100 mg/kg, p.o.	274.0 ± 3.50
<i>Ficus elasticleaves</i>	100 mg/kg p.o.	147.0 ± 4.81***
<i>Ficus elasticleaves</i>	200 mg/kg p.o.	123.0 ± 1.50***
Silymarin	2.5 mg/kg p.o.	113.0 ± 4.50***

**Table 9. Effect of *Ficus elastic* leaves and Silymarin on % ALP levels in Isoniazid induced hepatotoxicity in rats**

Treatment	Dose	ALP (%)
Normal	1 ml/kg, p.o.	162.0 ± 3.20
Isoniazid	100 mg/kg, p.o.	340.0 ± 5.57
<i>Ficus elasticleaves</i>	100 mg/kg p.o.	215.0 ± 4.30***
<i>Ficus elasticleaves</i>	200 mg/kg p.o.	178.0 ± 3.78***
Silymarin	2.5 mg/kg p.o.	167.0 ± 5.40***

**Table 10. Effect of Hydroalcoholic extract of *Ficus lyrataleaves* and Silymarin on % SGOT levels in Paracetamol induced hepatotoxicity in rats**

Treatment	Dose	SGOT (%)
Normal	1 ml/kg, p.o.	158 ± 2.5
Paracetamol	50 mg/kg, p.o.	352.45 ± 6.5
<i>Ficus lyrataleaves</i>	100 mg/kg p.o.	227.0 ± 3.5***
<i>Ficus lyrataleaves</i>	200 mg/kg p.o.	203.0 ± 3.0***
Silymarin	2.5 mg/kg p.o.	179.0 ± 3.5***

**Table 11. Effect of Hydroalcoholic extract of *Ficus lyrataleaves* and Silymarin on %SGPT levels in Paracetamol induced hepatotoxicity in rats**

Treatment	Dose	SGPT (%)
Normal	1 ml/kg, p.o.	151.0 ± 2.0
Paracetamol	50 mg/kg, p.o.	316.0 ± 3.50
<i>Ficus lyrataleaves</i>	100 mg/kg p.o.	189.0 ± 4.0***
<i>Ficus lyrataleaves</i>	200 mg/kg p.o.	174.0 ± 3.50***
Silymarin	2.5 mg/kg p.o.	152.0 ± 3.50***

**Table 12. Effect of *Ficus lyrataleaves* and Silymarin on % serum bilirubin levels in Paracetamol induced hepatotoxicity in rats**

Treatment	Dose	Serum Bilirubin (%)
Normal	1 ml/kg, p.o.	122.0 ± 5.50
Paracetamol	50 mg/kg, p.o.	304.0 ± 3.50
<i>Ficus lyrataleaves</i>	100 mg/kg p.o.	201.0 ± 4.51***
<i>Ficus lyrataleaves</i>	200 mg/kg p.o.	165.0 ± 1.60***
Silymarin	2.5 mg/kg p.o.	125.0 ± 4.50***

**Table 13. Effect of *Ficus lyrata* leaves and Silymarin on % ALP levels in Paracetamol induced hepatotoxicity in rats**

Treatment	Dose	ALP (%)
Normal	1 ml/kg, p.o.	155.0 ± 3.20
Paracetamol	50 mg/kg, p.o.	337.0 ± 5.57
<i>Ficus lyrata</i> leaves	100 mg/kg p.o.	232.0 ± 4.30***
<i>Ficus lyrata</i> leaves	200 mg/kg p.o.	191.0 ± 3.78***
Silymarin	2.5 mg/kg p.o.	175.0 ± 5.40***

**Table 14. Effect of Hydroalcoholic extract of *Ficus elastica* leaves and Silymarin on % SGOT levels in Paracetamol induced hepatotoxicity in rats**

Treatment	Dose	SGOT (%)
Normal	1 ml/kg, p.o.	165 ± 2.5
Paracetamol	50 mg/kg, p.o.	339.45 ± 6.5
<i>Ficus elastica</i> leaves	100 mg/kg p.o.	219.0 ± 3.5***
<i>Ficus elastica</i> leaves	200 mg/kg p.o.	181.0 ± 3.5***
Silymarin	2.5 mg/kg p.o.	168.0 ± 2.5***

**Table 15. Effect of Hydroalcoholic extract of *Ficus elastica* leaves and Silymarin on %SGPT levels in Paracetamol induced hepatotoxicity in rats**

Treatment	Dose	SGPT (%)
Normal	1 ml/kg, p.o.	152.0 ± 2.0
Paracetamol	50 mg/kg, p.o.	314.0 ± 3.50
<i>Ficus elastica</i> leaves	100 mg/kg p.o.	196.0 ± 4.0***
<i>Ficus elastica</i> leaves	200 mg/kg p.o.	174.0 ± 3.50***
Silymarin	2.5 mg/kg p.o.	156.0 ± 3.50***

**Table 16. Effect of *Ficus elastica* leaves and Silymarin on % serum bilirubin levels in Paracetamol induced hepatotoxicity in rats**

Treatment	Dose	Serum Bilirubin (%)
Normal	1 ml/kg, p.o.	120.0 ± 5.50
Paracetamol	50 mg/kg, p.o.	290.0 ± 3.50
<i>Ficus elastica</i> leaves	100 mg/kg p.o.	155.0 ± 4.81***
<i>Ficus elastica</i> leaves	200 mg/kg p.o.	131.0 ± 1.50***
Silymarin	2.5 mg/kg p.o.	125.0 ± 4.50***

**Table 17. Effect of *Ficus elastica* leaves and Silymarin on % ALP levels in Paracetamol induced hepatotoxicity in rats**

Treatment	Dose	ALP (%)
Normal	1 ml/kg, p.o.	162.0 ± 3.20
Paracetamol	50 mg/kg, p.o.	331.0 ± 5.57
<i>Ficus elastica</i> leaves	100 mg/kg p.o.	227.0 ± 4.30***
<i>Ficus elastica</i> leaves	200 mg/kg p.o.	172.0 ± 3.78***
Silymarin	2.5 mg/kg p.o.	165.0 ± 5.40***

### 3.4 Hepatoprotective Activity of *Ficus elastic* Leaves Extract Isoniazid Induced Hepatotoxicity

*In –Vivo* hepatoprotective activity of hydroalcoholic extract of *Ficus elastic* leaves and Silymarin on % SGOT levels, %SGPT levels, % serum bilirubin levels, % ALP levels in Isoniazid induced hepatotoxicity in rats.

### 3.5 Hepatoprotective Activity of *Ficus lyrata* Leaves Extract Paracetamol Induced Hepatotoxicity Model

Hepatoprotective activity of extract of *Ficus lyrata* leaves and Paracetamol on % SGOT levels, %SGPT levels, % serum bilirubin levels, % ALP levels in Paracetamol induced hepatotoxicity in rats. Tables 9,10,11,12. results of Hepatoprotective activity of hydroalcoholic extract of *Ficus elastic* leaves and Silymarin on % SGOT levels in Paracetamol induced hepatotoxicity in rats Tables 13,14,15,16.

### 3.6 Hepatoprotective Activity of *Ficus elastic* Leaves Extract Paracetamol Induced Hepatotoxicity Model

*In–Vivo* hepatoprotective activity of hydroalcoholic extract of *Ficus elastic* leaves and Silymarin on % SGOT levels, %SGPT levels, % serum bilirubin levels, % ALP levels in paracetamol induced hepatotoxicity in rats.

## 4. CONCLUSION

Hepatic damage is the main obstacle to drug development and a major factor in the reason why pharmaceutical companies have stopped selling their products. Liver ailment caused by medication might be either exceptional (low frequency and probably dosage related) or unsurprising (high rate and measurement related) [15,16] Responses that are eccentric, often referred to as unusual, can be classified as non-immune or resistive to extreme touchiness. Since ancient times, people have turned to plants as a source for healing illnesses, providing long-lasting excellent health, relieving severe suffering, and adding flavor and fragrance to food. India and China have been using a human services framework centered on plants for more than 5,000 years. Up until about 50 years ago, botanicals were used as common medications in Europe. Ancient advances from China, India, and

Greece were independently incorporated into the pharmacological systems of the Arabic countries [17,18].

However, the Ayurvedic framework is the most unparalleled in terms of study breadth profundity. This framework, which is said to have started approximately 6000 BC, is one of the oldest prescription frameworks. In many countries, medicinal plants have become an integral element of traditional medical frameworks. Documentation of the examination work done on conventional medications is required. These investigations aid in the identification of plant material. When synthetic analgesics and antimicrobial drugs were not yet widely available in the middle of the 20th century, natural drugs dominated the therapeutic landscape. People began to switch to this framework when allopathic pharmaceutical arrangements for manufactured drugs gained rapid traction and faster symptomatic relief [19].

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

Animal Ethic committee approval has been collected and preserved by the author(s).

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Handa SS. Future trends of plants as drugs. *Pharma Times*.1991;4:13-23.
2. Khandelwal KR. Ed. Practical Pharmacognosy Technique and Experiments, 23<sup>rd</sup>Edn: 2005; 15.
3. Kokate CK. Ed. Practical Pharmacognosy, 4<sup>th</sup>Edn., Vallabh Prakashan: 1994;112:120.
4. Kulikova O. Hepatoprotective Activity of Plant Peptides, *Ame. J. of Plant Sci*. 2015;6:848-55.



5. Levy SB, Marshall B. Antibacterial resistance worldwide: causes, challenges and responses. *Nature Medicine Supplement*. 2007;10(12):122-129.
6. Gonthier MP, Remesy C, Scalbert A, Cheynier V, Souquet JM, Poutanen K, Aura AM, Microbial metabolism of caffeic acid and its esters chlorogenic and caftaric acids by human faecal microbiota in vitro, *Biomedicine & Pharmacotherapy*. 2006; 60:536-540.
7. Mcintyre N, Benhamou JP, Bircher J, Rizzetto M, Rodes J. *Oxfordtext book of clinical hepatology*. Vol 2. 2nd ed. New York: Oxford University Press. 1991;907-09.
8. Mohamed ZM, Salem1 AZM, Salem 2\*, LM. Camacho3 and Hayssam M. Ali Antimicrobial activities and phytochemical composition of extracts of Ficus species: An over view, *African Journal of Microbiology Research*. 4211-4219.
9. Mohd Mujeeb, Shah Alam Khan, Vidhu Aeri, and Babar Ali, Hepatoprotective Activity of the Ethanolic Extract of *Ficus carica* Linn. Leavesin Carbon Tetrachloride-Induced Hepatotoxicity in Rats, *Iran J Pharm Res*. 2011;10(2): 301–306.
10. Nadeem M. et al. Hepatoprotective activity of some herbal formulations in India. *Indian Drugs*. 1996;33(8):390-395.
11. OECD, Guideline for Testing of Chemicals-Acute Oral Toxicity-Acute Toxic Class Method. Paris: OECD; 2001.
12. Patel R. BJD, Raval. Hepatoprotective effect of Indian Medicinal Palnts–Part I. *Indian Drugs*. 1998;25(6):224-27.
13. Reddy KN, Subbaraju GV, Reddy CS, Raju VS. Ethnoveterinary medicine for treating livestock in Eastern Ghats of Andhra Pradesh. *Indian Journal of Tradition Knowledge*. 2013;5(3):368-378.
14. Deepak Kumar, Mr. Dhyanendra Singh, Mrs. Pranali Mishra, Dr. Vishalgupta pharmacological evaluation of Hepatoprotective activity of fruits Extract of *Syzygium aeoranthum*. 2456-4184.
15. Anggun Selvy Arsyad, Arief Nurrochmad, Universitas Gadjah Mada, Nanang Fakhrudin Phytochemistry, traditional uses, and pharmacological activities of *Ficus elastica* Roxb. ex Hornem: A review. *Journal of Herbmед Pharmacology*. 2022;12(1):41-53. DOI:10.34172/jhp.2023.04
16. Hassan Abdalla Almahyl, Mawardi Rahmani%, Mohd Aspollah Sukar P & Abdul Manaf Ali4, Investigation on the Chemical Constituents of the Leaves of *Ficus elastica* Roxb. and Their Antimicrobial Activity, *Pertanika J. Sci. & Technol*. 2003;11(1):57–63.
17. Jean Emmanuel Mbosso Teinkela Xavier Siwe Noundou Edwige Laure Nguemfo Franck Meyer Rene Wintjens Michelle Isaacs Albert Emmanuel Mpondo Mpondo Heinrich C. Hoppe Rui Werner Maçedo Krause and Anatole Guy Blaise Azebaze Biological activities of plant extracts from *Ficus elastica* and *Selaginella vogelli*: An antimalarial, antitrypanosomal and cytotoxicity evaluation. *Saudi J Biol Sci*. 2018;25(1):117–122. Published online 2017 Jul 20. DOI: 10.1016/j.sjbs.2017.07.002
18. Katzung G. *Basicand Clinical Pharmacology*, 6thedi., Appleton and Lange, Connecticut. 2005;297-9.
19. Keppler D, Lesch R, Reutter W, Decker K. Experimental hepatitis induced by d-galactosamine. *Exp Mol Pathol*. 1968; 9(2):279-90.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/124456>