

Asian Research Journal of Gynaecology and Obstetrics

Volume 8, Issue 4, Page 24-31, 2022; Article no.ARJGO.93302

The Influence of Nutrition on Quality of Semen and Stamina Case Study: Vitolize and Bee Pollen Nutritional Products

Jackson Venuusto Modi Lado^{a*} and Zainab Hilal Hassan^b

^a Biotechnology Department, School of Applied and Industrial Sciences, University of Juba, Juba, South Sudan. ^b Centre for Nutritional Treatment, Fox LTD, Juba, South Sudan.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/93302

Original Research Article

Received: 23/10/2022 Accepted: 29/12/2022 Published: 31/12/2022

ABSTRACT

Several researches have focused on male infertility as an essential concern of man-life discontinuity. Roughly, 30-80% of infertility cases are because of oxidative stress and declined level of seminal total antioxidant capacity. There is few information about the influence of nutrition on quality of semen. The objective of this study was to detect the effects of nutritional products Vitolize and Bee pollen, which contain vitamin C and the B vitamins and other supplements for improving major semen parameters such as sperm concentration, motility, and morphology and fertility rate. A semen sample was obtained from a 42 years a male donor facing fertility problems. Analysed before and after treatment with the products in Queens laboratory, Nemra Talata. The results of this study indicated that after treatment, there was a significantly increase in sperm count, and an increase in progressive motility from 58.428 to 70.53 million/ejaculate, and 7.17 to14.29 % respectively. There was an increase in the number of live sperm cells, from 3 to 7 sperms. The

Asian Res. J. Gynaecol. Obst., vol. 8, no. 4, pp. 24-31, 2022

^{*}Corresponding author: E-mail: davidbilly772 @gmail.com;

current study also indicated that the products supplements for Vitolize and Bee pollen especially a combination of antioxidants such as vitamin C, vitamin E, and other supplements intake can effectively improve semen parameters in infertile men.

Keywords: Influence; vitolize; bee pollen; nutrition; quality; semen.

1. INTRODUCTION

"Barrenness can be defined as not being able to get the women into pregnancy state despite having continues, unprotected sex for at least 12 months" [1]. "More than 70 million couples facing infertility globally. Male infertility is an essential concern throughout the world. 14 % of couples suffer from barenness, based on research" [2]. "Male factors account for at least 50% of all infertility cases worldwide" [3]. Some factors such as radiation, smoking, varicocele, infection, urinary tract infection, environmental factors, nutritional deficiencies and oxidative stress as well as genetic factors can also be one of the reason for infertility. Due to Chromosome defects, chromosome abnormalities are found in which there is Deletion or duplication of whole chromosomes and inversion, or duplication of a portion of а chromosome or there is translocation of part of a chromosome to another chromosome. All these may contribute to male infertility [4,5]. "Oxidative stress occurs when the production of reactive oxygen species (ROS) exceeds the body's natural antioxidant defences" [6].

"These increased level of ROS can be resulted from environmental factors such as high temperature, electromagnetic waves, air pollution, insecticides, alcohol consumption, obesity and poor nutrition" [7]. There are evidence that sperms are simply affected by ROS and oxidative stress.

There are several researches that assist the role of ROS in male infertility theory [8]. "Products such as Vitolize and Bee pollen can introduce a high improvement for quality of semen, Vitolize and Bee pollen are some nutritional products, made for sperm improvement because they contain essential supplements such as Vitamins C and E. Ordinary antioxidants in semen include vitamin E, vitamin C, superoxide dismutase, glutathione and thioredoxin. These antioxidants neutralize free radical activity and protect sperm from ROS that already produced" [6].

2. MATERIALS AND METHODS

2.1 Study Area

The research was done in Queens Specialist private hospital, Queens Specialist private hospital is located around kokora road Juba, in Nimara Talata South Sudan. The hospital has simpler team to assist it giving faster, more useful care to patients and their families with an enhanced workflow [9].

2.2 Sample Collection

The sample was collected after 30 days and 4 days of abstinence, before and after treatment respectively from a 42 years adult, and then was taken to the laboratory for analysis.

2.3 Treatment prescription

The doses were applied as follows:

For Vitolize product, $1 \times 2/30$ days (one piece is taken in morning and one is taken in the evening for 30 days) and for Bee pollen, two are taken in morning and one is taken in the evening/for 30 days.

2.4 Sample Analysis

Semen samples were collected at the hospital by masturbation into a sterile plastic container. The samples were liquefied for 30 min in 37 °C before analysis. Macroscopic examination of semen was performed according to the 5th edition of WHO laboratory manual for the examination and processing of human semen [10]. Microscopic measurements of the sperm count, concentration, motility and morphology were determined with the use of computer-aided semen analysis (CASA). The basic components of the system were a bright field microscope (Olympus CX41, Tokyo, Japan), a digital camera to capture images (Olympus U-CMAD3), and a computer with software installed (SCA®Microptic S.L., Barcelona, Spain). The WHO [10] cut-off points were used to evaluate abnormal values of

semen quality parameters (Table 1). All analyses were performed by an experienced technician.nc.

2.5 Data Analysis

The computer also was equipped with Word Excel to analyse the data and compute the percentages from frequencies.

3. RESULTS

3.1 Physical Properties

The results for physical properties before and after the treatment showed that there was an increase for both sperm volume and pH, from 1.8 to 4 ml and 7.5 to 8 respectively (Table 1). Which are the good indicators for sperm improvement.

3.2 Results for Semen Analysis, Sperm Count and Progressive Motility

The results for semen analysis is illustrated in Table 2.

The results for semen analysis showed that after treatment there was an increase in sperm count, and progressive motility from 58.428 to 70.53 million/ejaculate, and 7.17 to14.29 % respectively (Table 2) and (Fig. 1).

3.3 Results for Progressive Motility

Results for progressive motility is represented in Fig. 1.

3.4 Results for Semen Analysis, the Dead and Live Sperm Estimation

The results for semen analysis, the dead and live sperm estimation is represented in Fig. 2.

The results showed that there was an increase in the number of live sperm cells, from 3 to 7 sperms (Fig. 2).

After treatment **Before treatment** Standard range Property Abstinence 30 Davs 4 Days 2-5 Days Collection Masturbation at lab Masturbation at lab Volume 1.8 ml 4 ml >=1.5 ml PH 7.5 8 >=7.2 Colour **Grey Opalescent Grey Opalescent** Viscosity Moderate Moderate Liquefaction time 30 Min 50 Min 15 - 60 min. **Liquefaction State** Complete Complete




Fig. 1. Progressive motility

Table 1. Sperm physical properties before and after treatment with vitolize nutritional product

Test	Before treatment with nutritional product Results	After treatment with nutritional product results	Status	Standard range
Concentration	32.46	23.51	passed	>= 15 million/ml
Total sperm count	58.428	70.53	passed	>=39 million/ejaculate
Progressive motility(PR)	7.17	14.29	failed	>=32%
Rapid PR	0.00	0.0		
Slow PR	7.17	14.29		
Total motility (PR+NP)	22.88	19.05	failed	>=40%
Vitality	60	87.5	passed	>=58%

Table 2. Test results for semen analysis, sperm count and progressive motility




Fig. 2. Semen analysis, the dead and live sperm estimation

4. DISCUSSION

It is believed that oxidative stress significantly affects male infertility. The results of this indicated that after treatment there was an increase in sperm count, and progressive motility from 58.428 to 70.53 million/ejaculate, and 7.17 to14.29 % respectively (Table 2) and (Fig. 1). And that there was an increase in the number of live sperm cells, from 3 to 7 sperms (Fig. 2).

"The results are all in line with other previous studies that focused on the effect of nutrition on sperm key parameters, their results showed a positive relationship between nutrition and semen parameters such as sperm count, motility and morphology" [11-13]. "One study indicated no significant relationship between nutrition therapy and semen parameters according to the smaller sample size compared to other researches" [14]. "A meta-analysis by Lafuente indicated that treatment with coenzyme Q10 led to a significant improvement in the sperm motility and density, whereas no significant improvements was observed in live birth and pregnancy rates [8]. The combination of vitamin E and vitamin C indicated no improvement in either sperm count or motility, but reduced sperm DNA damage" [15].

"Several studies have indicated that combined and N-acetvl-cvsteine selenium therapy improved male infertility" [2]. "Moslemi found that combination therapy with selenium and vitamin E significantly improved sperm motility and morphology. Two studies indicated that no improvement in sperm motility and morphology after zinc therapy" [16,17]. "However, two other studies showed a significant improvement in zinc-attached ligands and semen key enzymes after zinc therapy" [18,19].

Nutritional factor	Ма	ijor outcomes	References
Vitamin A		Normal blood-testis barrier function;	[20]
	-	Avoids germ-cell aplasia;	
	-	Fertile men have higher serum concentrations than infertile.	
Vitamin C	-	Improved sperm cell count, motility, and morphology;	
	-	Lower levels of vitamin C in seminal plasma of infertile men.	
	-	Higher live-birth rate, and a trend of better results of in vitro fertilization parameters;	[23-25]
	-	Decreases the lipid peroxidation of the sperm cell and seminal plasma;	
	-	Improves sperm cell motility;	
	-	Lower levels were found in men with oligozoospermia and asthenozoospermia.	
Vitamin D	-	The expression of vitamin D receptors and metabolizing enzymes are marked in human testis, ejaculatory tract,	[25,12]
		and mature sperm cells;	
	-	Positive association between serum levels and sperm motility;	
Vitamin B9		Protects against DNA damage.	[26,27]
Selenium	-	Protects against ROS;	[28,29]
	-	Deficiency promotes sperm cell abnormalities, and affects motility and fertility;	
Zinc	-	Important for spermatogenesis: cofactor of enzymes involved in DNA transcription and protein synthesis;	[30,18]
	-	Lower zinc levels in the seminal plasma of infertile men;	
	-	Increased the normal sperm cell morphology, sperm motility, and semen volume.	
N-acetylcysteine	-	Improved the volume, motility, and viscosity of sperm cells;	[29,31]
	-	Increased the serum total antioxidant capacity;	
	-	Reduced the serum peroxide and oxidative stress;	
	-	Increased sperm cell concentration, motility, and percent normal morphology in infertile men.	

"All studies which tested the influence of multiple antioxidants in a supplementation showed an improvement in semen parameters after therapy" [32,13]. "For example, combination therapy with carnitine, CoQ10, vitamin E and vitamin C for three to six months improved sperm concentration" [33]. "Only one study indicated a significant improvement in sperm concentration after combination therapy without enhancement in motility and morphology" [34].

5. CONCLUSION

In conclusion, in spite of several of reviewed researches indicated significant relation between antioxidant supplementations and one or two semen parameters but application of supplementations such as L-carnitine, selenium, vitamin C and vitamin E may contribute to enhancing sperm concentration, motility and morphology.

6. RECOMMENDATIONS

In order to maintain and increase the sperm vitality, mobility, concentration and mobility, the nutritional products such as Vitolize and Bee pollen should be taken regularly, at least once every six month because they contain most of the following (Table 3).

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Leaver RB. Male infertility: An overview of causes and treatment options. Br. J. Nurs. 2016;25:35–40.
- Irvine DS. Epidemiology and aetiology of male infertility; Hum Reprod 13 (suppl.). 1998:33-44.
- 3. Sharlip ID, Jarow JP, Belker AM, Lipshultz LI, Sigman M, Thomas AJ. Best practice policies for male infertility. Fertil steril. 2002;77: 873-882.
- Olayemi F. Review on some causes of male infertility .Afr J Biotech. 2010;9.

- Wong WY, Thomas CM, Merkus JM, Zielhuis GA, Steegers-Theunissen RP. Male factor subfertility: Possible causes and the impact of nutritional factors. Fertil Steril. 2000;73:435-442.
- Tremellen K. Oxidative stress and male infertility-a clinical perspective. Hum Reprod Update 2008;14:243-258.
- Aitken RJ, De Iuliis GN. Origins and consequences of DNA damage in male germ cells. Reprod Biomed Online. 2007;14: 727-733.
- Lafuente R, González-Comadrán M, Solà I, López G, Brassesco M, Carreras R et al. Coenzyme Q10 and male infertility: a meta-analysis. J Assis Reprod Gen. 2013; 30:1147-1156.
- 9. Queens-medical-centre-South.Sudan.
- World Health Organization. WHO Laboratory Manual for the Examination and Processing of Human Semen, 5th Ed.; WHO Press: Geneva, Switzerland; 2010. ISBN 978-924-154-778-9.
- 11. Radigue C, Es-Slami S, Soufir J. Relationship of carnitine transport across the epididymis to blood carnitine and androgens in rats. Arch Androl 1996;37: 27-31.
- Blomberg Jensen M, Bjerrum PJ, Jessen TE, Nielsen JE, Joensen UN, Olesen IA, Petersen JH, Juul A, Dissing S, Jørgensen N. Vitamin D is positively associated with sperm motility and increases intracellular calcium in human spermatozoa. Hum. Reprod. 2011;26:1307–1317.
- Gopinath P, Kalra B, Saxena A, Malik S, Kochhar K, Kalra S et al. Fixed dose combination therapy of antioxidants in treatment of idiopathic oligoasthenozoospermia: Results of a randomized, double-blind, placebocontrolled Clinical Trial. Int J Infertil Fetal Med. 2013;4:6-13.
- 14. Balercia G, Regoli F, Armeni T, Koverech A, Mantero F, Boscaro M. Placebocontrolled doubleblind randomized trial on the use of L-carnitine, Lacetylcarnitine, or combined L-carnitine and Lacetylcarnitine in men with idiopathic asthenozoospermia. Fertil Steril. 2005;84: 662-671.
- Gvozdjáková A, Kucharská J, Dubravicky J, Mojto V, Singh RB. Coenzyme Q10, α-Tocopherol, and Oxidative Stress Could

Be Important Metabolic Biomarkers of Male Infertility. Dis mark 2015;2015: ID 827941.

- Ebisch I, Thomas C, Peters W, Braat D, SteegersTheunissen R. The importance of folate, zinc and antioxidants in the pathogenesis and prevention of subfertility. Hum Reprod Update. 2007; 13:163-174.
- Hadwan MH, Almashhedy LA, Alsalman ARS. Study of the effects of oral zinc supplementation on peroxynitrite levels, arginase activity and NO synthase activity in seminal plasma of Iraqi asthenospermic patients. Reprod Biol Endocrin. 2014;12:1.
- 18. Colagar AH, Marzony ET, Chaichi MJ. Zinc levels in seminal plasma are associated with sperm quality in fertile and infertile men. Nutr. Res. 2009;29: 82–88.
- 19. Ebisch I, Pierik F, De Jong F, Thomas C, Steegers-Theunissen R. Does folic acid and zinc sulphate intervention affect endocrine parameters and sperm characteristics in men? Int J Androl. 2006; 29:339-345.
- Al-Azemi M, Omu A, Fatinikun T, Mannazhath N, Abraham S. Factors contributing to gender differences in serum retinol and α-tocopherol in infertile couples. Reprod. Biomed. Online. 2009;19:583– 590.
- Colagar AH, Marzony ET. Ascorbic acid in human seminal plasma: Determination and Its relationship to sperm quality. J. Clin. Biochem. Nutr. 2009;45:144–149.
- 22. Akmal M, Qadri JQ, Al-Waili NS, Thangal S, Haq A, Saloom KY. Improvement in human semen quality after oral supplementation of vitamin C. J. Med. Food. 2006;9:440–442.
- Al-Azemi M, Omu A, Fatinikun T, Mannazhath N, Abraham S. Factors contributing to gender differences in serum retinol and α-tocopherol in infertile couples. Reprod. Biomed. Online. 2009;19:583–590.
- 24. Matorras R, Pérez-Sanz J, Corcóstegui B, Pérez-Ruiz I, Malaina I, Quevedo S, Aspichueta F, Crisol L, Martinez-Indart L, Prieto B. Effect of vitamin E administered to men in infertile couples on sperm and assisted reproduction outcomes: A doubleblind randomized study. F&S Rep. 2020;1:219–226.

- Blomberg Jensen M, Nielsen JE, Jorgensen A, Rajpert-De Meyts E, Kristensen DM, Jorgensen N, Skakkebaek NE, Juul A, Leffers H. Vitamin D receptor and vitamin D metabolizing enzymes are expressed in the human male reproductive tract. Hum. Reprod. 2010;25: 1303–1311.
- 26. Hoek J, Steegers-Theunissen RPM, Willemsen SP, Schoenmakers S. Paternal folate status and sperm quality, pregnancy outcomes, and epigenetics: A systematic review and meta-analysis. Mol. Nutr. Food Res. 2020;64:1900696.
- Boonyarangkul A, Vinayanuvattikhun N, Chiamchanya C, Visutakul P. Comparative study of the effects of tamoxifen citrate and folate on semen quality of the infertile male with semen abnormality. J. Med. Assoc. Thai. 2015;98:1057–1063.
- Ahsan U, Kamran Z, Raza I, Ahmad S, Babar W, Riaz MH, Iqbal Z. Role of selenium in male reproduction—A review. Anim. Reprod. Sci. 2014;146:55–62.
- Safarinejad MR, Safarinejad S. Efficacy of selenium and/or N-Acetyl-Cysteine for improving semen parameters in infertile men: A double-blind, placebo controlled, randomized study. J. Urol. 2009;181:741– 751.
- Zhao J, Dong X, Hu X, Long Z, Wang L, Liu Q, Sun B, Wang Q, Wu Q, Li L. Zinc levels in seminal plasma and their correlation with male infertility: A systematic review and meta-analysis. Sci. Rep. 2016;6:22386.
- Ciftci H, Verit A, Savas M, Yeni E, Erel O. Effects of N-acetylcysteine on semen parameters and oxidative/antioxidant status. Urology. 2009;74:73–76.
- 32. Galatioto GP, Gravina GL, Angelozzi G, Sacchetti A, Innominato PF, Pace G et al. May antioxidant therapy improve sperm parameters of men with persistent oligospermia after retrograde embolization for varicocele? World J Urol. 2008;26:97-102.
- Gvozdjáková A, Kucharská J, Dubravicky J, Mojto V, Singh RB. Coenzyme Q10, α-tocopherol, and oxidative stress could be important metabolic biomarkers of male infertility. Dis mark 2015; 2015: ID 827941.

Lado and Hassan; Asian Res. J. Gynaecol. Obst., vol. 8, no. 4, pp. 24-31, 2022; Article no.ARJGO.93302

34. Safarinejad MR, Safarinejad S. Efficacy of selenium and/or N-acetyl-cysteine for improving semen parameters in infertile

men: a double-blind, placebo controlled, randomized study. J Urol 2009;181: 741-751.

© 2022 Lado and Hassan; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

> Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/93302