



Economic Analysis of Bell Pepper (*Capsicum annuum* L.) Production as Influenced by Different Organic Nutrient Sources under Mid Hill Condition of Himachal Pradesh

Atal Hameedi ^{a*}, Kuldeep Singh Thakur ^b, Nesar Ahmad Nesar ^c, Meenakshi Attri ^c, Rahmat Gul Stanikzai ^c, Rakesh Kumar ^a and Satesh Kumar ^a

^a Division of Vegetable Science & Floriculture, Faculty of Agriculture, Sher-e-Kashmir University of Agricultural Science and Technology, Jammu, Main Campus, Chatha-180009, Jammu and Kashmir, India.

^b Department of Vegetable Science, Dr. Y.S. Parmar University of Horticulture and Forestry, Solan-173230 (H.P.), India.

^c Division of Agronomy, Faculty of Agriculture, Sher-e-Kashmir University of Agricultural Science and Technology, Jammu, Main Campus, Chatha-180009, Jammu and Kashmir, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i1030792

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/86588>

Received 15 February 2022

Accepted 28 April 2022

Published 05 May 2022

Original Research Article

ABSTRACT

A field investigation was conducted during the Kharif of 2016 at the Department of Vegetable Science, UHF, Nauni, Solan to explore the economic analysis of bell pepper (*Capsicum annuum* L.) production as influenced by organic nutrient sources. The experiment was designed in Randomized Complete Block Design (RCBD) Factorial with three replications comprising 10 different treatments. The study indicated that vermicompost @ 7 t/ha + Jeevamrut (Drenching @ 5% + Foliar spray @ 3%) highly affected yield and yield attributing traits of bell pepper and documented higher fruit weight (59.33 g), no. of fruits/plant (29.13), fruit yield/plot (24.73 kg), and fruit yield/ha (366.42 q) along with the highest gross income (₹ 916,050.00/ha) and net return (₹ 713,795.00/ha) whereas the greatest B: C ratio (5.43) was attained with no organic manure +

Jeevamrut (Drenching @ 5% + Foliar spray @ 3%). As a result, from an economic standpoint, Jeevamrut (Drenching @ 5% + Foliar spray @ 3%) application and Vermicompost @ 7 t/ha in combination with Jeevamrut (Drenching @ 5% + Foliar spray @ 3%) application for high yield may be suggested for commercial production.

Keywords: Organic manure; liquid manure; economics; yield; bell pepper.

1. INTRODUCTION

Bell pepper (*Capsicum annuum* L. var. *grossum* Sendt.) is an important member of Solanaceae family, having chromosome number $2n = 24$, and recognized as sweet pepper, capsicum, and Shimla mirch. It is a Mexican native, with a secondary origin in Guatemala [1]. It is the world's third most popular vegetable crop after tomatoes, having thick flesh and low-calorie content despite being rich in nutrients, especially vitamin A and vitamin C. [2]. Organic farming has the ability to ameliorate soil productivity, biodiversity, and economic return [3]. Organic agriculture is gaining popularity, and organic food markets are fast developing in numerous nations, comprising India [4]. Organic amendments might be used as a substitute for inorganic fertilizers [5]. The excessive use of inorganic fertilizers caused series of problems among which soil degradation, poor quality production, environmental pollution and some serious hazards to human health are the most common. Application of chemical fertilizers can only provide one or two essential elements. Organic manure plays an important function in plant growth by providing all needed macro elements and microelements in conveniently accessible forms in the course of mineralization [6], and soil physical and chemical qualities are improved [7]. Manures can therefore produce a high yield, a high gross revenue, and a high net return.

2. MATERIALS AND METHODS

The current investigation was carried out at the Vegetable Research Farm of the Department of Vegetable Science, UHF, Nauni, Solan, HP from April to September 2016. The experiment was conducted in a Randomized Complete Block Design (RCBD) Factorial with three replications including 10 treatment combinations of manure and liquid manure, namely; T_1 (M_0L_0): No organic manure + No liquid manure (Control), T_2 (M_1L_0): FYM @ 20 t/ha + No liquid manure, T_3 (M_2L_0) : Vermicompost @ 7 t/ha + No liquid manure, T_4

(M_3L_0): FYM @ 10 t/ha + VC 3.5 t/ha + No liquid manure, T_5 (M_4L_0): FYM @ 15 t/ha + VC 1.75 t/ha + No liquid manure, T_6 (M_0L_1) : No organic manure + Jeevamrut (Drenching + Foliar Spray), T_7 (M_1L_1): FYM @ 20 t/ha. + Jeevamrut (Drenching+ Foliar Spray), T_8 (M_2L_1): Vermicompost @ 7 t/ha. + Jeevamrut (Drenching + Foliar Spray), T_9 (M_3L_1): FYM @ 10 t/ha + VC 3.5 t/ha + Jeevamrut (Drenching + Foliar Spray), T_{10} (M_4L_1): FYM @ 15 t/ha + VC 1.75 t/ha + Jeevamrut (Drenching + Foliar Spray). On March 5, 2016, seeds of bell pepper cv. Solan Bharpur has been sown in nursery beds, and seedlings have been transplanted on April 26, 2016. The plot was 240 cm x 225 cm in size, with a spacing of 0.6 m X 0.45 m, and benefit: cost (BC) ratios were recorded. As indicated by, the data collected on different factors were examined for RBD design [8]. The data were interpreted using the 'F' test value, and the crucial difference (CD) was determined at a 5% level of significance. Before planting, the soil was analyzed and found to be rich in organic matter having OC 1.38 per cent, neutral in reaction (pH 7.2), normal in EC (0.431dSm^{-1}) medium in available nitrogen (395.14 kg/ha) and potassium (263.2 kg/ha) and high in phosphorus (58.24 kg/ha).

2.1 Application of organics

Before transplanting of bell pepper seedlings, all FYM and Vermicompost were applied properly by blending them with soil as per treatment combination per plot, and liquid manure has been applied as 3 drenching with Jeevamrut @ 5% at 15 day intervals started at the time of transplanting + 2 Foliar Spray of Jeevamrut @ 3% at 15 day intervals started after 45 days of transplanting.

2.2 Disease and Pest Management

To inhibit the growth of soil pathogenic fungi, 5 kg of *Trichoderma viride* (powder formulation) was mixed with 100 kg of FYM and covered with polythene for 7 days. Before putting it in the field, the mixture was mixed every 3-4 days. On bell

pepper plants, 2-3 cc of *Verticillium lecanii* (liquid formulation) was sprayed to suppress aphids.

2.3 Climate

The area's climate is mainly sub-temperate and semi-humid, with chilly winters. In general, the warmest months are May and June, while the coldest are December and January. The yearly rainfall fluctuates between 1000 and 1300 mm, the majority of which is falling at the time of monsoon season (June-September). The average temperature ranged from 21.5 to 24.4 degree Celsius, with relative humidity ranging from 41% to 83% during the crop season.

2.4 Economics of Treatments

The economics of treatment are pivotal factors to examine before recommending it to a farmer. The cost of bell pepper production under different treatments was investigated in order to determine the economics of various organic nutrient sources while taking into account the current price of inputs and produce. The gross return was calculated using the market price of the produce. The following equations were used to calculate net returns and benefit-cost ratios for each nutritional treatment:

Net return (₹/ha) = Gross returns (₹/ha) - Cost of cultivation (₹/ha)

Benefit Cost ratio: = Net return (₹/ha) /Cost of cultivation (₹/ha)

3. RESULT AND DISCUSSION

The attentive study of the data in Table 1 displayed that the organic and liquid manure fertilization had a significant impact on bell pepper yield and yield contributing traits individually as well as their combined effect. Among different levels of manure, vermicompost treated plots achieved significantly more fruits per plant (27.00), average fruit weight (56.43 g), fruit breadth (5.69 cm), fruit length (6.54 cm), fruit yield (21.77 kg/ plot), and fruit yield (322.47 q/ha), while no manure (M₀) produced the least.

In comparison to no Jeevamrut (L₀), Jeevamrut (L₁) produced the highest number of fruits per plant (25.75), average fruit weight (54.43 g), fruit width (5.62 cm), fruit yield

(20.20 kg/plot), and fruit yield (299.26 q/ha). The effect of Jeevamrut on fruit length (cm), as well as their combined effect with different organic manure levels, observed to be non-significant. The combined effect of both the factors revealed that M₂L₁ had the highest number of fruits (29.13/plant), average fruit weight (59.33 g), fruit width (5.89 cm) fruit yield (24.73 kg/ plot), and fruit yield (366.42 q/ha), whereas M₀L₀ had the lowest of all these characteristics.

Jeevamrut may have contributed to increased vegetative growth, the quantity of fruits per plant, increased nutrient distribution during the critical growth phase, and increased carbohydrate assimilation and translocation. It might be because of the use of vermicompost, which has a direct influence on plant development by providing a stable provenance of plant macro and microelements. However, some of these essential elements exist in readily available forms to the plants, the maximum of which are liberated gently and depositing minerals, resulting a proper supply of nutrients all the time to the plant by dynamic and consistent source of essential elements.

It might also be attributable to improved soil physical, chemical, and biological qualities, as well as supplementation of macro and micro nutrients, as a result of the combined impact of vermicompost and Jeevamrut, which improved growth, yield, and yield contributing attributes. These findings are also corroborated by [9,10,11,12,13,14,15,16] found that Vermicompost had a favorable impact on pepper.

3.1 Economics of the Treatment

The economics of the various treatment combinations depicted in Table 2. A keen study of data divulged that highest cost of production Rs. 202,255.00 was incurred in M₂L₁, whereas lowest (₹ 94,030.00) was observed in M₀L₀. Similarly maximum gross income per hectare amounting to ₹ 916,050.00 was obtained in M₂L₁ followed by M₁L₁ (₹ 787,650.00), while minimum gross income (₹ 502,225.00) was recorded in M₀L₀. This is due to the high cost of production and maximum marketable yield in M₂L₁ as compared to other treatments used in the study. However, highest net return (₹ 713,795.00) was recorded in M₂L₁ followed by M₁L₁ (₹ 652,695.00) and lowest net return (₹ 408,195.00) was observed in M₀L₀.

Table 1. Yield and yield contributing traits as influenced by different organic nutrient sources [9]

Particular	Numberof fruits per plant	Fruit length (cm)	Fruit breadth(cm)	Fruit weight (g)	Fruit yieldkg/plot	Fruityieldq/ha
Organic manure						
M ₀	20.63	5.90	5.25	45.27	15.13	224.15
M ₁	25.23	6.33	5.62	55.00	19.98	296.05
M ₂	27.00	6.54	5.69	56.43	21.77	322.47
M ₃	24.87	6.51	5.50	52.37	19.22	284.69
M ₄	24.37	6.29	5.39	50.13	18.17	269.14
CD _{0.05} (M)	0.64	0.24	0.03	1.30	0.41	6.04
Liquid Manure						
L ₀	23.09	6.25	5.36	49.25	17.51	259.34
L ₁	25.75	6.38	5.62	54.43	20.20	299.26
CD _{0.05} (L)	0.41	NS	0.02	0.82	0.26	3.82
Interaction (M×L)						
M ₀ L ₀	18.67	5.72	5.22	42.47	13.56	200.89
M ₀ L ₁	22.60	6.08	5.29	48.07	16.70	247.41
M ₁ L ₀	24.40	6.38	5.42	53.47	18.70	277.04
M ₁ L ₁	26.07	6.28	5.83	56.53	21.27	315.06
M ₂ L ₀	24.87	6.50	5.48	53.53	18.80	278.52
M ₂ L ₁	29.13	6.57	5.89	59.33	24.73	366.42
M ₃ L ₀	24.13	6.50	5.38	48.80	18.57	275.06
M ₃ L ₁	25.60	6.53	5.62	55.93	19.87	294.32
M ₄ L ₀	23.40	6.14	5.32	48.00	17.90	265.19
M ₄ L ₁	25.33	6.43	5.45	52.27	18.43	273.09
CD_{0.05} (M×L)	0.91	NS	0.04	1.84	0.58	8.54

Table 2. Economics of the bell pepper production as influenced by different treatments

Treatments	Fruit yieldq/ha	Gross income(₹/ha)	Total cost of production (₹/ha)	Net return(₹/ha)	Benefit CostRatio
M ₀ L ₀	200.89	502,225.00	94,030.00	408,195.00	4.34
M ₁ L ₀	277.04	692,600.00	132,830.00	559,770.00	4.21
M ₂ L ₀	278.52	696,300.00	200,130.00	496,170.00	2.48
M ₃ L ₀	275.06	687,650.00	166,480.00	521,170.00	3.13
M ₄ L ₀	265.19	662,975.00	149,655.00	513,320.00	3.43
M ₀ L ₁	247.41	618,525.00	96,155.00	522,370.00	5.43
M ₁ L ₁	315.06	787,650.00	134,955.00	652,695.00	4.84
M ₂ L ₁	366.42	916,050.00	202,255.00	713,795.00	3.53
M ₃ L ₁	294.32	735,800.00	168,605.00	567,195.00	3.36
M ₄ L ₁	273.09	682,725.00	151,780.00	530,945.00	3.50

In overall, maximum B: C ratio (5.43) was recorded in M₀L₁ followed by B: C ratio (4.84) in M₁L₁, whereas minimum B: C ratio (2.48) was calculated in M₂L₀, which may be due to the lower cost of production and comparable gross income in M₂L₁. Hence the application of M₀L₁ (No manure + Jeevamrut) rated as the most economic treatment for bell pepper production under mid hill conditions of Himachal Pradesh [17]

4. CONCLUSION

Based on the current research, it can be stated that the combination application of Vermicompost + Jeevamrut produced the greatest results for the majority of yield and yield contributing attributes. It also produced the greatest gross revenue (₹ 916,050.00/ha) and net return (₹ 713,795.00/ha), whereas MOL1 (No organic manure + Jeevamrut) produced the highest benefit cost ratio (5.43). As a consequence, based on the findings of the current research, the use of Jeevamrut (Drenching at 5% + Foliar spray at 3%), from an economic standpoint, may be suggested for organic bell pepper growing, as it resulted in the highest B: C ratio (5.43). In contrast, application of vermicompost at a rate of 7 t/ha in conjunction with Jeevamrut (Drenching at 5% + Foliar spray at 3%) is advised for increased yield and long-term crop production. Further, if the farmer produces vermicompost on farm, definitely it will reduce the cost of cultivation incurred for getting better yield and sustainable crop production & good economic return as well.

Authors have declared that no competing interests exist.

REFERENCES

1. Bukasov SM. The cultivated plants of Mexico, Guatemala and Columbia. Instituta Rastenievodostva Vaskhnil, Leningrad. 1930;553.
2. Raturi HC, Uppal GS, Singh SK, Kachwaya DS. Effect of organic and inorganic nutrient sources on growth, yield and quality of bell pepper (*Capsicum annuum* L.) grown under polyhouse condition. Journal of Pharmacognosy and Phytochemistry. 2019;8(1):1788-1792.
3. Hameedi A, Thakur KS, Sharma U, Yousafzai A, Mohammadi MH, Durrani H, Durani A. Effect of organic nutrient sources on NPK uptake, soil nutrient status and yield of bell pepper (*Capsicum annuum* L.) under mid hill condition of Himachal Pradesh. International Journal of Chemical Studies. 2018;6(1):1913-1917.
4. Gopinath KA, Saha S, Mina BL. Effects of organic amendments on productivity and profitability of bell pepper–french bean–garden pea system and on oil properties during transition to organic production. Communications in Soil Science and Plant Analysis. 2012; 42:2572-2585.
5. Naeem M, Iqbal J, Bakhsh MAA. Comparative study of inorganic fertilizers and organic manures on yield and yield components of Mungbean. Journal of Agriculture & Social Sciences, 2006;2(4):227-239.
6. Nweke IA, Ijearu SI, Igili DN. Effect of different sources of animal wastes on the growth and yield of okra (*Abelmoschus esculentus* L.Moench) in Ustoxic Dystropept at Enugu South Eastern, Nigeria. International Journal of Scientific and Technology Research. 2013;2(3): 135-137.
7. Chaterjee B, Ghanti P, Thapa U, Tripathy P. Effect of organic nutrition in sprouting broccoli. Vegetable Science. 2005;33(1):51-54.
8. Gomez AA, Gomez AA. Statistical Procedures for Agricultural Research. John Wiley and Sons. New York. 1984;680.
9. Arancon NQ, Edwards CA, Bierman P, James DM, Stephen L, and Christie W. Effects of vermicomposts on growth and marketable fruits of field-grown tomatoes, peppers and strawberries. Pedobiologia. 2003;47:731-735.
10. Natesh N, Vyakaranahal BS, Shekhargouda M, Deshpande VK. Effect of micronutrients and organics on growth, seed yield and quality of chilli. Karnataka Journal Agricultural Sciences. 2005;18(2):334-337.
11. Arancon NQ, Edward CA, Biermanb P, Metzgerc JD, Lucht C. Effects of vermicompost produced from cattle manure, food waste and paper waste on the growth and yield of peppers in the field. Pedobiologia. 2005;49:297-306.
12. Joshi R, Pal Vig A. Effect of vermicompost on growth, yield and

- quality of tomato (*Lycopersicum esculentum* L). African Journal of Basic & Applied Sciences. 2010;2(3-4):117-123.
13. Kumar BM. Effect of vermicompost on germination, growth and yield of vegetable plants. Scrutiny International Research Journal of Agriculture, Plant Biotechnology and Bio Products. 2016;3(1):07-13.
 14. Ramesh G, Ajithkumar K, Savitha AS, Patil SG. Integrated influence of organic manures in addition to inorganic fertilizers on growth, yield parameters and early blight disease of tomato (*Lycopersicon esculentum* L.). International Journal of Biological & Pharmaceutical Research. 2015;6(6):478-483.
 15. Ahmed N, Rameshwar, Sini JP, Sharma RP, Punam, Seth M. Performance of chickpea under organic and inorganic sources of nutrients at different soil moisture regimes in chickpea-okra cropping system. Himachal Journal of Agricultural Research. 2017; 43: 23-28.
 16. Huerta E, Vidal O, Jarquin A, Geissen V, Gomez R. Effect of vermicompost on the growth and production of amashito pepper, interactions with earthworms and rhizobacteria. Compost Science & Utilization. 2010;18:282-288.
 17. Adhikari P, Khanal A, Subedi R. Effect of different sources of organic manure on growth and yield of sweet pepper. Advances in Plants & Agriculture Research. 2016;3:00111.

© 2022 Hameedi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/86588>