



Bluprins® as Alternative Bud Break Promoter for 'Maxi Gala' and 'Fuji Suprema' Apple Trees

**Cristhian Leonardo Fenili¹, José Luiz Petri², André Amarildo Sezerino²,
Mariuccia Schlichting De Martin², Gentil Carneiro Gabardo¹
and Eduardo da Silva Daniel^{1*}**

¹*Agroveterinary Sciences Center, Santa Catarina State University, Lages, Brazil.*

²*EPAGRI/Caçador Experimental Station, Caçador, Brazil.*

Authors' contributions

This work was carried out in collaboration between all authors. Author CLF performed the statistical analysis and wrote the first draft of the manuscript. Author JLP designed the study and wrote the protocol. Authors AAS and MSDM managed the analyses of the study. Authors GCG and ESD managed the literature searches and the translation of the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2018/43649

Editor(s):

- (1) Dr. Suleyman Korkut, Professor, Duzce University, Department of Forest Industrial Engineering, Division of Wood Mechanic and Technology, Turkey.
(2) Dr. Daniele De Wrachien, Professor, State University of Milan, Italy.

Reviewers:

- (1) Gariglio, Norberto Francisco, Universidad Nacional del Litoral, Argentina.
(2) George Nanos, University of Thessaly, Greece.
(3) Raúl Leonel Grijalva-Contreras, Instituto Nacional de Investigaciones Forestales Agrícolas y Pecuarias, México.

Complete Peer review History: <http://www.sciencedomain.org/review-history/26381>

Original Research Article

Received 23 June 2018
Accepted 06 September 2018
Published 24 September 2018

ABSTRACT

Aims: The effect of Bluprins® at different concentrations combined with calcium nitrate and ammonium nitrate on phenology, bud break induction and fruit production of 'Maxi Gala' and 'Fuji Suprema' apple tree cultivars was evaluated in mild winter conditions.

Study Design: The experiment was designed in randomised block and replicated five times.

Place and Duration of Study: The experiment was carried out in the municipality of Caçador, Brazil, during the growing seasons of 2013/2014, 2014/2015, 2015/2016 and 2016/2017.

Methodology: The study considered 'Maxi Gala' and 'Fuji Suprema' apple trees. The treatments evaluated were as follows: 1. Control (untreated); 2. Mineral oil 3.5% + hydrogen cyanamide 0.35%; 3. Bluprins® 3.0% + calcium nitrate 3.0%; 4. Bluprins® 5.0% + calcium nitrate 3.0%;

*Corresponding author: E-mail: edudaniel@hotmail.com;

5. Bluprins[®] 3.0% + calcium nitrate 5.0%; 6. Bluprins[®] 5.0% + calcium nitrate 5.0%; 7. Bluprins[®] 3.0% + calcium nitrate 3.0% + ammonium nitrate 3.0%; 8. Bluprins[®] 5.0% + calcium nitrate 4.0% + ammonium nitrate 4.0%. Phenology, axillary and terminal bud break, fruit set, fruit production and average fruit weight were evaluated. The phenological stage of green tip (C-C3) and the beginning of bud break and flowering were anticipated by the application of Bluprins[®] and hydrogen cyanamide in compared to the control. The axillary and terminal bud break was increased by the application of bud break promoters for both cultivars considering the four growing seasons studied. The average fruit weight did not show significant differences between treatments in the growing seasons of 2013/2014 and 2015/2016. Bluprins[®], in combination with calcium nitrate and ammonium nitrate, proved effective in inducing bud break of 'Maxi Gala' and 'Fuji Suprema' apple tree cultivars under mild winter conditions. The phenomenon is less effective with mineral oil in combination with hydrogen cyanamide. Bluprins[®] in combination with calcium nitrate and ammonium nitrate anticipates the bud break and flowering period and reduces the flowering period for 'Maxi Gala' and 'Fuji Suprema' apple tree cultivars in compared to plants without application of bud break promoters. Bluprins[®] does not affect considerably the fruit set and fruit production of 'Maxi Gala' and 'Fuji Suprema' apple tree cultivars.

Keywords: *Malus domestica* Borkh.; dormancy; mild winter chilling; bud break induction.

1. INTRODUCTION

The apple-growing (*Malus domestica* Borkh.) has a high production volume, estimated at 91.4 million tons of harvested fruit in 2016, which classifies apple as one of the most produced fruits in the world [1]. Apple production is of great importance in Southern Brazil. In this region, more than 4 thousand fruit growers harvested 1.05 million tons of apples in that year, representing a financial turnover of R\$ 6 billion and generating 195 thousand of jobs in marketing sector [2].

Apple trees present suspension of vegetative growth in winter, called dormancy. This mechanism allows it to survive periods of low temperatures [3]. To overcome dormancy, plants have to satisfy their chilling requirements to initiate spring bud break, shoot meristematic extension growth and anthesis [4]. Chilling requirements vary depending on the cultivar [5]. In Brazil, 90% of apple orchards are composed by 'Gala' and 'Fuji' cultivars and their spontaneous mutations [6]. These cultivars present chilling requirements of around 600 to 800 hours of temperatures below 7.2°C to overcome dormancy [7]. The insufficient chilling accumulation in a specified cultivar results mainly in the reduction of bud break and uneven flowering [8]. These dysfunctions have economic consequences due to the impact on fruit production and quality, may compromise management orchard techniques such as chemical thinning and influence the next harvest by reducing the production due to the lower

formation of reproductive structures in the plants [9,10].

In some subtropical climate countries such as Brazil, a few regions present favourable conditions to overcome apple tree dormancy [11,6]. In these areas, the solution for the cultivation has been the selection of cultivars with low chilling requirement combined with the application of bud break promoters and cultural practices to break bud dormancy. These proceedings provide adequate bud break and flowering [12,13].

The desirable characteristics of chemical substances have good efficiency for the bud break induction, low cost and minimum toxicity to plants and environment [14]. Despite the existence of a large number of effective substances for bud break induction, only a few are used commercially. The high cost of application and high toxicity of the compounds are the primary limiting factors [15]. Hydrogen cyanamide combined with mineral oil is the most effective compound for bud break induction and is extensively used in the cultivation of apple trees and other temperate climate fruit trees for more than 20 years in Brazil [16,17]. However, the toxicity of hydrogen cyanamide is diversifying the standard recommendation to break dormancy of apple trees by alternative substances which is less harmful to the agrochemical applicator and environment [18].

Organic nitrogen compounds have shown potential for commercial use. The combination of

these inducers with calcium nitrate has proved promising for bud break induction of apple trees [19,10]. Several studies carried out in different countries have shown promising results of Bluprins® bio-stimulant on bud break induction and flowering uniformity for trees of table grapes, cherries and kiwifruit. Preliminarily, positive results were also obtained for apple and peach trees [20]. Bluprins® is a concentrated gel formulation for breaking bud dormancy of temperate fruit trees containing polysaccharides, amino acids, nitrogen and organic carbon [21].

In this context, the study aimed to test the efficiency of Bluprins® combined with calcium nitrate and ammonium nitrate as an alternative to hydrogen cyanamide in relation to bud break induction in 'Maxi Gala' and 'Fuji Suprema' apple tree cultivars.

2. MATERIALS AND METHODS

The experiment was carried out in an experimental orchard located in the municipality of Caçador, Santa Catarina State, Brazil (26°50'S, 50°58'O, 941 m a.s.l). Eight-year-old plants of 'Maxi Gala' and 'Fuji Suprema' apple trees were grafted on rootstock Marubakaido and M-9 as interstock. The orchard density was about 2,500 plants ha⁻¹ and the plants were trained to a central leader system. Orchard management practices were applied according to the recommendations for apple production system [22].

The experiment was carried out in a randomised complete block design following a 4 x 8 factorial structure (four growing seasons and eight bud break promoters) with five replications of a single tree. The growing seasons evaluated were 2013/2014, 2014/2015, 2015/2016 and 2016/2017, and the bud breakers evaluated were: 1. Control (untreated); 2. Mineral oil 3.5% + hydrogen cyanamide 0.35%; 3. Bluprins® 3.0% + calcium nitrate [Ca(NO₃)₂] 3.0%; 4. Bluprins® 5.0% + calcium nitrate 3.0%; 5. Bluprins® 3.0% + calcium nitrate 5.0%; 6. Bluprins® 5.0% + calcium nitrate 5.0%; 7. Bluprins® 3.0% + calcium nitrate 3.0% + ammonium nitrate [NH₄(NO₃)] 3.0%; 8. Bluprins® 5.0% + calcium nitrate 4.0% + ammonium nitrate 4.0%. The bio-stimulant used is the commercial product Bluprins®, composed of water, ammonium nitrate, sugar cane molasses, amino acids, citric acid, sodium hydroxide, supplying 4% ammoniacal N, 4% nitric N, 0.7% organic N and 5.5% organic C. The commercial product Dormex®, has 52% of active

ingredient, used as a source of hydrogen cyanamide. The commercial product Assist® was used as a source of mineral oil (75.6%). Compounds were applied with a motorised backpack sprayer. Application time was performed on 06/09/2013, 03/09/2014, 26/08/2015 and 25/08/2016 for the growing seasons of 2013/2014, 2014/2015, 2015/2016 and 2016/2017, respectively. During the winter period, the growing seasons of 2013/2014, 2014/2015, 2015/2016 and 2016/2017 presented 544, 271, 189 and 612 hours of temperature below 7.2°C and 940, 884, 746 and 1,200 of chilling units (Modified North Carolina model [23]), respectively (Fig. 1).

The phenology, axillary and terminal bud break and fruit set were evaluated. The evaluation of flowering phenology consisted of determining the dates of occurrence of the green tip stage (C-C3), start of flowering, full bloom and end of flowering [22,18]. The C-C3 stage was considered when 50% of the dormant buds presented "green tip" (beginning of sprouting). The beginning of flowering was considered when the plants had 5% of the flowers in opening stage, full bloom more than 80% and the termination of flowering was defined when the last flowers were open. The axillary bud break was obtained by counting both burst and dormant buds in five one-year-old shoots previously selected, located in the middle third of the plant. A scaffold branch was selected to estimate the percentage of terminal bud break. These data were collected at 30 and 60 days after dormancy breaking (DADB). The fruit set was obtained as a percentage in 100 flower cluster in the same scaffold branch used to estimate terminal bud break. Fruit production per tree and average fruit weight was also measured and reported.

The data were statistically evaluated by using analysis of variance (ANOVA). Percentage data were transformed by the formula $\arcsin [(x + 1) / 100]^{1/2}$ before being subjected to ANOVA. Treatment means were compared by using the Scott-Knott test at 5% probability level. The statistical analysis was performed by the Sisvar program version 5.6 [24].

3. RESULTS AND DISCUSSION

For both 'Maxi Gala' and 'Fuji Suprema' cultivars, the treatments resulted in a significant difference ($p < 0.05$) for most of the evaluated variables. For 'Fuji Suprema' cultivar, there was a significant




Fig. 1. Historical minimum (Min.) and maximum (Max.) average temperatures (°C) and minimum and maximum temperatures (°C) of the growing seasons of 2013/2014, 2014/2015, 2015/2016 and 2016/2017 during the dormancy period of apple trees. Caçador, SC

interaction between growing seasons and bud breaker promoters considering axillary and terminal bud break. However, for 'Maxi Gala' cultivar, there was no significant interaction between growing seasons and bud breaker promoters for any of the evaluated variables (data not shown).

For 'Maxi Gala' apple trees, the phenological stage C-C3 was different between the control treatment and the other treatments. The application of mineral oil in combination with hydrogen cyanamide advanced this stage 19 days, 21 days, 24 days and 23 days compared to the control treatment plant in the growing seasons of 2013/2014, 2014/2015, 2015/2016 and 2016/2017, respectively. The treatments containing Bluprins[®] also advanced this stage from 5 to 15 days in 2013/2014, from 13 to 19 days in 2014/2015, from 5 to 7 days in 2015/2016 and from 14 to 17 days in comparison to the control treatment in 2016/2017 (Table 1).

For 'Fuji Suprema' apple trees, there was not a defined period that characterised the C-C3 stage for the control treatment. In the 2013/2014 growing season; considering the other treatments, this stage occurred practically at the same date. In 2014/2015 growing season, the treatments containing Blueprins[®] advanced the C-C3 stage from 1 to 8 days in relation to the control treatment and delayed this stage from 2 to 5 days in relation to mineral oil + hydroxygen cyanamide application. In 2015/2016 and 2016/2017 growing seasons, all the treatments advanced this stage from 2 to 8 days and from 4 to 10 days, respectively, in relation to the control treatment (Table 2).

The bud break was also advanced in both cultivars by the application of bud break promoters. However, there were small differences between mineral oil + hydrogen cyanamide and the Bluprins[®] treatments. The beginning, full bloom and termination of flowering were advanced in relation to the control treatment for 'Maxi Gala' and 'Fuji Suprema' cultivars in the four growing seasons, and the Bluprins[®] treatments showed a tendency to delay these phenological stages a few days in relation to treatment mineral oil + hydrogen cyanamide (Tables 1 and 2).

The growing seasons of 2014/2015 and 2015/2016 represented lower chilling accumulation and more extended flowering period than that of the growing seasons of 2013/2014 and 2016/2017. According to Kozmá et al. [25], duration of the flowering period is influenced by environmental conditions, being longer under low chilling accumulation during the winter season.

In 2013/2014 growing season, the flowering period was prolonged in the control treatment in compared to the other treatments. For 'Maxi Gala', the flowering period comprised of 14 days in the control trees, whereas the other treatments varied between 11 to 15 days. For 'Fuji Suprema', the flowering period comprised of 15 days in the control trees, and varied between 6 to 16 days in the other treatments. In 2014/2015, 2015/2016 and 2016/2017 growing seasons, the other treatments resulted in a shortened flowering period in compared to the control treatment only for 'Fuji Suprema', varying from 8 to 12 days, and from 10 to 17 days, respectively in 2015/2016 and 2016/2017, while the control

trees had a flowering period of 32 and 18 days, respectively. For 'Maxi Gala', the treatments presented this period equal to the control treatment or longer in the 2014/2015 and 2016/2017 growing seasons (Tables 1 and 2).

Petri and Leite [7] endorsed that prolonged flowering periods may difficult for some cultural practices such as thinning and disease control,

due to the occurrence of different phenological stages within the same plant. The duration of the flowering period can evaluate the efficiency of bud break promoters, and the most effective treatments are those with shorter flowering period and more uniform flowering, ripening and harvesting. It is revealed that Bluprins® acts in the advance and shortening of the flowering period of apple trees under mild winter conditions.

Table 1. Phenological stages of 'Maxi Gala' apple trees under the influence of compounds for bud break in four growing seasons (Date/Month). Caçador, SC, 2018

Treatments	C-C3	Bud break	Flowering		
			Start	Full bloom	End
2013/2014					
1. Control	10/10	14/10	14/10	21/10	28/10
2. MO 3,5% + HC 0,35%	21/09	06/10	08/10	12/10	19/10
3. B 3% + Ca(NO ₃) ₂ 3%	05/10	08/10	08/10	20/10	23/10
4. B 5% + Ca(NO ₃) ₂ 3%	25/09	08/10	08/10	18/10	23/10
5. B 3% + Ca(NO ₃) ₂ 5%	05/10	08/10	08/10	20/10	23/10
6. B 5% + Ca(NO ₃) ₂ 5%	25/09	06/10	08/10	12/10	23/10
7. B 3%+Ca(NO ₃) ₂ 3%+NH ₄ (NO ₃) 3%	25/09	06/10	08/10	18/10	25/10
8. B 5%+Ca(NO ₃) ₂ 4%+NH ₄ (NO ₃) 4%	05/10	12/10	12/10	21/10	26/10
2014/2015					
1. Control	12/10	12/10	12/10	24/10	28/10
2. MO 3,5% + HC 0,35%	21/09	25/09	28/09	03/10	14/10
3. B 3% + Ca(NO ₃) ₂ 3%	29/09	02/10	04/10	12/10	20/10
4. B 5% + Ca(NO ₃) ₂ 3%	26/09	29/09	29/09	12/10	18/10
5. B 3% + Ca(NO ₃) ₂ 5%	26/09	29/09	30/09	12/10	21/10
6. B 5% + Ca(NO ₃) ₂ 5%	23/09	25/09	30/09	08/10	18/10
7. B 3%+Ca(NO ₃) ₂ 3%+NH ₄ (NO ₃) 3%	28/09	01/10	01/10	12/10	21/10
8. B 5%+Ca(NO ₃) ₂ 4%+NH ₄ (NO ₃) 4%	29/09	03/10	04/10	12/10	21/10
2015/2016					
1. Control	10/10	-	10/10	28/10	05/11
2. MO 3,5% + HC 0,35%	16/09	21/09	21/09	24/09	30/09
3. B 3% + Ca(NO ₃) ₂ 3%	23/09	25/09	27/09	08/10	26/10
4. B 5% + Ca(NO ₃) ₂ 3%	23/09	25/09	25/09	01/10	04/10
5. B 3% + Ca(NO ₃) ₂ 5%	21/09	22/09	25/09	28/09	04/10
6. B 5% + Ca(NO ₃) ₂ 5%	21/09	25/09	26/09	11/10	26/10
7. B 3%+Ca(NO ₃) ₂ 3%+NH ₄ (NO ₃) 3%	21/09	24/09	25/09	30/09	09/10
8. B 5%+Ca(NO ₃) ₂ 4%+NH ₄ (NO ₃) 4%	21/09	22/09	25/09	28/09	04/10
2016/2017					
1. Control	05/10	08/10	08/10	16/10	20/10
2. MO 3,5% + HC 0,35%	12/09	17/09	21/09	30/09	06/10
3. B 3% + Ca(NO ₃) ₂ 3%	21/09	30/09	30/09	08/10	10/10
4. B 5% + Ca(NO ₃) ₂ 3%	19/09	30/09	30/09	06/10	10/10
5. B 3% + Ca(NO ₃) ₂ 5%	21/09	30/09	30/09	06/10	10/10
6. B 5% + Ca(NO ₃) ₂ 5%	21/09	30/09	30/09	05/10	15/10
7. B 3%+Ca(NO ₃) ₂ 3%+NH ₄ (NO ₃) 3%	18/09	22/09	30/09	06/10	10/10
8. B 5%+Ca(NO ₃) ₂ 4%+NH ₄ (NO ₃) 4%	18/09	20/09	25/09	21/10	08/10

MO: Mineral oil; HC: Hydrogen cyanamide; B: Bluprins®

Table 2. Phenological stages of 'Fuji Suprema' apple trees under the influence of compounds for bud break in four growing seasons (Date/Month). Caçador, SC, 2018

Treatments	C-C3	Bud break	Flowering		
			Start	Full bloom	End
2013/2014					
1. Control	-	07/10	07/10	15/10	22/10
2. MO 3,5% + HC 0,35%	25/09	04/10	05/10	08/10	16/10
3. B 3% + Ca(NO ₃) ₂ 3%	25/09	06/10	06/10	15/10	22/10
4. B 5% + Ca(NO ₃) ₂ 3%	25/09	03/10	05/10	08/10	15/10
5. B 3% + Ca(NO ₃) ₂ 5%	25/09	04/10	05/10	08/10	16/10
6. B 5% + Ca(NO ₃) ₂ 5%	25/09	04/10	05/10	08/10	16/10
7. B 3%+Ca(NO ₃) ₂ 3%+NH ₄ (NO ₃) 3%	25/09	03/10	06/10	08/10	12/10
8. B 5%+Ca(NO ₃) ₂ 4%+NH ₄ (NO ₃) 4%	23/09	04/10	03/10	07/10	15/10
2014/2015					
1. Control	01/10	03/10	10/10	20/10	28/10
2. MO 3,5% + HC 0,35%	25/09	27/09	30/09	04/10	08/10
3. B 3% + Ca(NO ₃) ₂ 3%	30/09	02/10	06/10	10/10	18/10
4. B 5% + Ca(NO ₃) ₂ 3%	28/09	30/09	01/10	08/10	12/10
5. B 3% + Ca(NO ₃) ₂ 5%	28/09	01/10	03/10	10/10	18/10
6. B 5% + Ca(NO ₃) ₂ 5%	29/09	30/09	03/10	10/10	15/10
7. B 3%+Ca(NO ₃) ₂ 3%+NH ₄ (NO ₃) 3%	23/09	29/09	28/09	03/10	18/10
8. B 5%+Ca(NO ₃) ₂ 4%+NH ₄ (NO ₃) 4%	27/09	30/09	03/10	10/10	18/10
2015/2016					
1. Control	25/09	25/09	26/09	-	28/10
2. MO 3,5% + HC 0,35%	17/09	20/09	18/09	25/09	30/09
3. B 3% + Ca(NO ₃) ₂ 3%	23/09	23/09	22/09	27/09	30/09
4. B 5% + Ca(NO ₃) ₂ 3%	20/09	22/09	21/09	26/09	30/09
5. B 3% + Ca(NO ₃) ₂ 5%	19/09	21/09	21/09	26/09	30/09
6. B 5% + Ca(NO ₃) ₂ 5%	17/09	21/09	18/09	26/09	30/09
7. B 3%+Ca(NO ₃) ₂ 3%+NH ₄ (NO ₃) 3%	17/09	21/09	19/09	26/09	30/09
8. B 5%+Ca(NO ₃) ₂ 4%+NH ₄ (NO ₃) 4%	19/09	21/09	21/09	26/09	30/09
2016/2017					
1. Control	24/09	03/10	26/09	06/10	14/10
2. MO 3,5% + HC 0,35%	14/09	23/09	20/09	30/09	07/10
3. B 3% + Ca(NO ₃) ₂ 3%	20/09	26/09	26/09	04/10	12/10
4. B 5% + Ca(NO ₃) ₂ 3%	18/09	23/09	25/09	30/09	07/10
5. B 3% + Ca(NO ₃) ₂ 5%	14/09	25/09	25/09	01/10	05/10
6. B 5% + Ca(NO ₃) ₂ 5%	14/09	24/09	20/09	30/09	05/10
7. B 3%+Ca(NO ₃) ₂ 3%+NH ₄ (NO ₃) 3%	14/09	20/09	20/09	30/09	04/10
8. B 5%+Ca(NO ₃) ₂ 4%+NH ₄ (NO ₃) 4%	15/09	25/09	25/09	01/10	07/10

MO: Mineral oil; HC: Hydrogen cyanamide; B: Bluprins®

The bud break promoters maximized the axillary bud break at 30 and 60 DADB for 'Maxi Gala' and 'Fuji Suprema' apple trees in the four growing seasons (Tables 3 and 4). According to Petri [26], the rate of axillary bud break is the variable that better expresses the effectiveness of bud break promoters, and can be used as indicative of cultivar adaptation to local environmental conditions. However, the efficiency of bud break promoters depends, in addition to the cultivar, on the vigour of the plant, time of application and concentration of the bud break promoter.

For 'Maxi Gala' cultivar, the treatment mineral oil + hydrogen cyanamide showed higher axillary bud break in relation to the other treatments at 30 and 60 DADB in the four growing seasons. The treatments Bluprins® 3% + Ca(NO₃)₂ 3% and Bluprins® 5% + Ca(NO₃)₂ 3% differed from the other treatments in the 2013/2014 growing season. In 2014/2015 and 2016/2017, there were no significant differences between Bluprins® treatments and the control treatment. In 2015/2016, the treatment Bluprins® 3% + Ca(NO₃)₂ 5% provided higher axillary bud break in relation to the control treatment and other Bluprins® treatments (Table 3).

For 'Fuji Suprema' apple trees at 30 and 60 DADB, all treatments showed higher axillary bud break compared to the control treatment in the 2013/2014 growing season, without differences among them. Pasa et al. [10] reported that a nutritive solution containing calcium nitrate and mineral oil showed similar kind of effects to hydrogen cyanamide on axillary bud break of this cultivar. In 2014/2015, except for the treatment Bluprins® 3% + Ca(NO₃)₂ 3%, all treatments presented higher axillary bud break compared to the control treatment, for this cultivar. The treatment mineral oil + hydrogen cyanamide presented the highest axillary bud break, followed by Bluprins® 5% + Ca(NO₃)₂ 3%. In 2015/2016, the treatment Bluprins® 5% + Ca(NO₃)₂ 4% + NH₄(NO₃) 4% was superior to the other Bluprins® treatments at 30 DADB. At 60 DADB, this treatment and Bluprins® 3% + Ca(NO₃)₂ 3% + NH₄(NO₃) 3% treatment were superior to the other Bluprins® treatments. However, both treatments were lower than the treatment of mineral oil + hydrogen cyanamide at 30 and 60 DADB. In 2016/2017, the treatment Bluprins® 5% + Ca(NO₃)₂ 4% + NH₄(NO₃) 4% was superior to the other Bluprins® treatments at 30 and 60 DADB, and did not differ from the treatment mineral oil + hydrogen cyanamide (Table 4).

Hawerroth et al. [27] reported lower axillary bud break in 'Imperial Gala' than in 'Fuji Suprema' cultivar, corroborating the higher difficulty in inducing bud break in 'Gala' cultivars. The low

axillary bud break, similar to that verified in this work, has already been endorsed by Leite et al. [28], conclude that temperate fruit trees cultivated in subtropical climate conditions, where the chilling requirement is not satisfied, represent low bud break levels associated with high bud break and flowering heterogeneity along the branches.

For 'Maxi Gala' apple trees, considering terminal bud break, the treatment mineral oil + hydrogen cyanamide showed higher values than that of other treatments at 30 and 60 DADB in the 2013/2014 and 2015/2016 growing seasons. This treatment and treatment of Bluprins® 3% + Ca(NO₃)₂ 3%, achieved higher terminal bud break than the other treatments at 30 DADB in 2014/2015 and at 60 DADB in 2016/2017. The Bluprins® treatments were superior to the control at 30 DADB in 2013/2014, 2014/2015 and 2015/2016. In 2016/2017, all the Bluprins® treatments were superior to the control at 30 and 60 DADB, and at 30 DADB, they did not differ from the treatment mineral oil + hydrogen cyanamide (Table 5). Marchi et al. [29] reported that even terminal buds, required low stimulus to break the dormancy, [30] only showed a high bud break by the application of mineral oil + hydrogen cyanamide. However, Pasa et al. [10] endorsed no significant differences between plants treated with hydrogen cyanamide, a nutrient solution containing calcium nitrate and control (untreated) plants, considering terminal bud break.

Table 3. Axillary bud break (%) of 'Maxi Gala' apple trees under the influence of compounds for bud break during four growing seasons. Caçador, SC, 2018

Treatments	Axillary bud break (%)							
	2013/2014		2014/2015		2015/2016		2016/2017	
	30 DADB	60 DADB	30 DADB	60 DADB	30 DADB	60 DADB	30 DADB	60 DADB
1. Control	1.1 c	3.8 c	1.2 b	4.9 b	0.9 c	1.8 c	0.0 b	1.0 b
2. MO 3,5% + HC 0,35%	41.5 a	43.8 a	20.9 a	26.0 a	23.0 a	25.7 a	13.5 a	24.4 a
3. B 3% + Ca(NO ₃) ₂ 3%	22.0 b	23.7 b	2.5 b	9.5 b	2.2 c	4.6 c	0.9 b	2.0 b
4. B 5% + Ca(NO ₃) ₂ 3%	22.4 b	25.3 b	3.9 b	7.4 b	1.8 c	1.8 c	3.1 b	3.8 b
5. B 3% + Ca(NO ₃) ₂ 5%	9.2 c	12.5 c	3.8 b	10.0 b	8.5 b	9.8 b	5.7 b	7.7 b
6. B 5% + Ca(NO ₃) ₂ 5%	16.6 c	19.6 c	6.7 b	10.2 b	3.1 c	4.7 c	3.5 b	5.3 b
7. B 3%+Ca(NO ₃) ₂ 3% +NH ₄ (NO ₃) 3%	11.8 c	16.5 c	3.8 b	5.9 b	1.6 c	3.5 c	3.5 b	7.6 b
8. B 5%+Ca(NO ₃) ₂ 4% +NH ₄ (NO ₃) 4%	8.2 c	9.8 c	5.1 b	10.7 b	2.2 c	3.1 c	3.5 b	4.9 b
CV (%)	72.2	61.4	55.8	33.2	66.2	57.6	69.0	43.0

MO: Mineral oil; HC: Hydrogen cyanamide; B: Bluprins®. DADB: Days after dormancy breaking; CV: coefficient of variation. Means followed by same letter do not differ by Scott-Knott test at 5% probability

Table 4. Axillary bud break (%) of 'Fuji Suprema' apple trees under the influence of compounds for bud break during four growing seasons. Caçador, SC, 2018

Treatments	Axillary bud break (%)							
	2013/2014		2014/2015		2015/2016		2016/2017	
	30 DADB	60 DADB	30 DADB	60 DADB	30 DADB	60 DADB	30 DADB	60 DADB
1. Control	3.3 b	12.6 b	0.0 d	5.2 d	0.0 d	4.4 c	15.2 c	22.6 c
2. MO 3,5% + HC 0,35%	53.1 a	60.2 a	48.0 a	51.9 a	50.1 a	50.1 a	65.5 a	81.0 a
3. B 3% + Ca(NO ₃) ₂ 3%	39.3 a	43.0 a	2.6 d	7.5 d	2.9 d	5.7 c	20.2 c	25.8 c
4. B 5% + Ca(NO ₃) ₂ 3%	52.7 a	59.4 a	22.3 b	30.3 b	5.1 c	7.3 c	32.3 b	48.4 b
5. B 3% + Ca(NO ₃) ₂ 5%	48.0 a	58.5 a	14.4 c	20.3 c	1.6 d	7.2 c	41.2 b	45.9 b
6. B 5% + Ca(NO ₃) ₂ 5%	56.1 a	62.3 a	11.3 c	14.2 c	0.6 d	4.6 c	33.4 b	37.2 b
7. B 3%+Ca(NO ₃) ₂ 3% +NH ₄ (NO ₃) 3%	72.4 a	73.8 a	13.7 c	20.0 c	7.9 c	11.0 b	25.2 c	29.2 c
8. B 5%+Ca(NO ₃) ₂ 4% +NH ₄ (NO ₃) 4%	55.6 a	71.3 a	11.4 c	14.9 c	18.8 b	19.9 b	67.3 a	70.9 a
CV (%)	27.8	27.9	47.2	31.0	49.0	35.4	25.5	21.5

MO: Mineral oil; HC: Hydrogen cyanamide; B: Bluprins[®]. DADB: Days after dormancy breaking; CV: coefficient of variation. Means followed by same letter do not differ by Scott-Knott test at 5% probability

For 'Fuji Suprema' apple trees, all the treatments showed higher terminal bud break compared to the control at 30 DADB in 2013/2014. At 60 DADB, the treatments Bluprins[®] 3% + Ca(NO₃)₂ 3% and Bluprins[®] 3% + Ca(NO₃)₂ 5% provided lower terminal bud break than the other Bluprins[®] treatments and the treatment mineral oil + hydrogen cyanamide. The treatment mineral oil + hydrogen cyanamide was superior to the other treatments in 2014/2015 at 30 DADB, whereas the treatment Bluprins[®] 3% + Ca(NO₃)₂ 3% was superior to the control treatment and other Bluprins[®] treatments. At 60 DADB, the treatments mineral oil + hydrogen cyanamide,

Bluprins[®] 3% + Ca(NO₃)₂ 3% and Bluprins[®] 5% + Ca(NO₃)₂ 3% were superior to the other treatments. In 2015/2016, at 60 DADB, the treatments Bluprins[®] 3% + Ca(NO₃)₂ 5%, Bluprins[®] 5% + Ca(NO₃)₂ 3% and Bluprins[®] 5% + Ca(NO₃)₂ 4% + NH₄(NO₃) 4% did not differ from the control, while the other treatments were superior to the control, not differing from the treatment mineral oil + hydrogen cyanamide. In 2016/2017, only the treatment Bluprins[®] 5% + Ca(NO₃)₂ 3% did not differ from the control treatment. The other treatments presented higher terminal bud break and did not differ from each other at 30 and 60 DADB (Table 6).

Table 5. Terminal bud break (%) of 'Maxi Gala' apple trees under the influence of compounds for bud break during four growing seasons. Caçador, SC, 2018

Treatments	Terminal bud break (%)							
	2013/2014		2014/2015		2015/2016		2016/2017	
	30 DADB	60 DADB	30 DADB	60 DADB	30 DADB	60 DADB	30 DADB	60 DADB
1. Control	15.9 c	60.3 b	12.6 c	52.9 ^{ns}	8.2 c	35.6 b	26.7 b	45.7 c
2. MO 3,5% + HC 0,35%	96.1 a	96.4 a	71.8 a	77.1	80.6 a	90.8 a	77.2 a	90.6 a
3. B 3% + Ca(NO ₃) ₂ 3%	57.8 b	57.5 b	43.9 b	68.1	27.9 b	49.2 b	55.4 a	66.8 b
4. B 5% + Ca(NO ₃) ₂ 3%	46.8 b	68.4 b	36.0 b	59.2	37.6 b	61.2 b	53.4 a	70.1 b
5. B 3% + Ca(NO ₃) ₂ 5%	42.2 b	60.3 b	32.0 b	72.1	38.4 b	60.8 b	59.5 a	83.5 a
6. B 5% + Ca(NO ₃) ₂ 5%	68.3 b	78.5 b	45.9 b	63.2	34.5 b	65.1 b	57.7 a	69.6 b
7. B 3%+Ca(NO ₃) ₂ 3% +NH ₄ (NO ₃) 3%	62.9 b	71.9 b	51.3 b	73.3	33.4 b	44.0 b	63.1 a	72.8 b
8. B 5%+Ca(NO ₃) ₂ 4% +NH ₄ (NO ₃) 4%	51.0 b	64.8 b	37.3 b	71.7	29.6 b	56.7 b	65.4 a	74.2 b
CV (%)	24.8	17.9	29.8	18.3	26.1	21.9	24.2	15.6

MO: Mineral oil; HC: Hydrogen cyanamide; B: Bluprins[®]. DADB: Days after dormancy breaking; CV: coefficient of variation. Means followed by same letter do not differ by Scott-Knott test at 5% probability. ns: not significant

Table 6. Terminal bud break (%) of 'Fuji Suprema' apple trees under the influence of compounds for bud break during four growing seasons. Caçador, SC, 2018

Treatments	Terminal bud break (%)							
	2013/2014		2014/2015		2015/2016		2016/2017	
	30 DADB	60 DADB	30 DADB	60 DADB	30 DADB	60 DADB	30 DADB	60 DADB
1. Control	46.7 c	80.2 b	24.3 c	80.3 b	30.9 ^{ns}	83.6 b	69.2 b	82.6 b
2. MO 3,5% + HC 0,35%	89.7 a	94.0 a	86.3 a	96.2 a	73.3	99.4 a	95.0 a	97.7 a
3. B 3% + Ca(NO ₃) ₂ 3%	68.5 b	86.2 b	59.2 b	87.5 a	52.5	98.3 a	78.5 b	91.3 b
4. B 5% + Ca(NO ₃) ₂ 3%	88.9 a	100 a	44.1 c	88.2 a	53.1	97.0 a	95.1 a	97.8 a
5. B 3% + Ca(NO ₃) ₂ 5%	74.5 b	87.9 b	44.9 c	77.8 b	48.4	93.6 b	95.3 a	95.8 a
6. B 5% + Ca(NO ₃) ₂ 5%	95.2 a	97.8 a	29.2 c	82.7 b	46.1	92.3 b	91.0 a	97.5 a
7. B 3%+Ca(NO ₃) ₂ 3% +NH ₄ (NO ₃) 3%	96.3 a	99.6 a	37.8 c	70.0 b	63.2	95.6 a	92.6 a	98.2 a
8. B 5%+Ca(NO ₃) ₂ 4% +NH ₄ (NO ₃) 4%	90.7 a	99.4 a	36.0 c	81.2 b	59.9	80.3 b	96.2 a	99.3 a
CV (%)	16.0	7.1	28.1	14.1	21.9	12.2	13.1	10.7

MO: Mineral oil; HC: Hydrogen cyanamide; B: Bluprins[®]. DADB: Days after dormancy breaking; CV: coefficient of variation. Means followed by same letter do not differ by Scott-Knott test at 5% probability. ^{ns}: not significant

For both 'Maxi Gala' and 'Fuji Suprema' apple trees, the high terminal bud break confirms the excellent efficiency of Bluprins[®] associated with calcium nitrate regardless of its concentration.

For 'Maxi Gala' apple trees, considering fruit set, the treatments Bluprins[®] 3% + Ca(NO₃)₂ 3% and Bluprins[®] 3% + Ca(NO₃)₂ 3% + NH₄(NO₃) 3% were superior to the other treatments in the 2015/2016 growing season; in 2013/2014, 2014/2015 and 2016/2017, the treatments showed no significant differences (Table 7). For 'Fuji Suprema' apple trees and fruit set, the treatments Bluprins[®] 5% + Ca(NO₃)₂ 3% and Bluprins[®] 3% + Ca(NO₃)₂ 3% + NH₄(NO₃) 3% did not differ from the control treatment and were significantly superior to the other treatments, in 2013/2014; in 2014/2015 and 2015/2016, the treatments showed no differences, and in 2016/2017, the control treatment showed a higher fruit set in relation to the other treatments (Table 8). El-Agamy et al. [31] endorsed a negative effect of the treatments with hydrogen cyanamide on the fruit set for Anna cultivar. Erez [14], Petri and Leite [7] argued the possibility of reduction in the fruit set when bud break promoters are applied due to non-synchronisation of the pollination between cultivars under conditions of insufficient chilling accumulation during the winter period. The fruit set may also be reduced due to weather conditions that affect the activity of pollinators and the pollen viability [32]. However, the high fruit set values obtained in some treatments

indicates that there were no problems related to pollination and that the concentration of the flowering period for the treatments with bud break promoters did not reduce the fruit set, even though the flowering period was more concentrated in the treatments with bud break promoters in comparison to the control treatment. According to Erez [14], the use of bud break promoters may result in a drastic reduction of the fruit set due to the nutritional competition established between vegetative and reproductive sinks. The Bluprins[®] treatments were equal or superior to the treatment of mineral oil + hydrogen cyanamide for fruit set of both cultivars in all the growing seasons.

For 'Maxi Gala' apple trees, considering the fruit production per tree, the treatments Bluprins[®] 5% + Ca(NO₃)₂ 3%, Bluprins[®] 3% + Ca(NO₃)₂ 3% + NH₄(NO₃) 3% and Bluprins[®] 5% + Ca(NO₃)₂ 4% + NH₄(NO₃) 4% resulted in higher values compared to the other treatments, in the 2013/2014 growing season. Apple trees treated with Bluprins[®] 3% + Ca(NO₃)₂ 3% + NH₄(NO₃) 3% produced 19.9 kg tree⁻¹ and the control treatment, 9.0 kg tree⁻¹, an increase of 121.1%. There were no significant differences between treatments in the 2014/2015 growing season (Table 9). The harvesting was not evaluated in 2015/2016. In 2016/2017 growing season, the treatments Bluprins[®] 3% + Ca(NO₃)₂ 3% + NH₄(NO₃) 3% and mineral oil + cyanamide hydrogen resulted in higher fruit production per tree than the other treatments.

Table 7. Fruit set (%) of 'Maxi Gala' apple trees under the influence of compounds for bud break during four growing seasons. Caçador, SC, 2018

Treatments	Fruit set (%)			
	2013/2014	2014/2015	2015/2016	2016/2017
1. Control	16.8 ^{ns}	11.7 ^{ns}	12.8 b	2.2 ^{ns}
2. MO 3,5% + HC 0,35%	21.8	9.5	6.9 b	1.0
3. B 3% + Ca(NO ₃) ₂ 3%	23.4	10.4	86.7 a	37.5
4. B 5% + Ca(NO ₃) ₂ 3%	30.2	13.4	43.3 b	23.3
5. B 3% + Ca(NO ₃) ₂ 5%	17.7	4.8	25.7 b	4.7
6. B 5% + Ca(NO ₃) ₂ 5%	20.9	29.6	3.9 b	1.5
7. B 3%+Ca(NO ₃) ₂ 3%+NH ₄ (NO ₃) 3%	20.1	10.6	140.7 a	11.2
8. B 5%+Ca(NO ₃) ₂ 4%+NH ₄ (NO ₃) 4%	28.0	15.6	40.0 b	10.4
CV (%)	69.2	51.8	102.1	161.0

MO: Mineral oil; HC: Hydrogen cyanamide; B: Bluprins[®]. CV: coefficient of variation. Means followed by same letter do not differ by Scott-Knott test at 5% probability. ^{ns}: not significant

Table 8. Fruit set (%) of 'Fuji Suprema' apple trees under the influence of compounds for bud break during four growing seasons. Caçador, SC, 2018

Treatments	Fruit Set (%)			
	2013/2014	2014/2015	2015/2016	2016/2017
1. Control	380.0 a	242.0 ^{ns}	159.3 ^{ns}	359.4 a
2. MO 3,5% + HC 0,35%	181.9 b	205.5	76.9	109.6 b
3. B 3% + Ca(NO ₃) ₂ 3%	260.2 b	266.3	95.6	208.7 b
4. B 5% + Ca(NO ₃) ₂ 3%	300.5 a	163.3	65.9	70.5 b
5. B 3% + Ca(NO ₃) ₂ 5%	192.8 b	257.9	49.9	191.8 b
6. B 5% + Ca(NO ₃) ₂ 5%	224.9 b	168.3	61.2	200.5 b
7. B 3%+Ca(NO ₃) ₂ 3%+NH ₄ (NO ₃) 3%	304.3 a	164.7	37.3	148.6 b
8. B 5%+Ca(NO ₃) ₂ 4%+NH ₄ (NO ₃) 4%	170.1 b	173.4	29.0	148.3 b
CV (%)	34.5	39.9	57.1	32.1

MO: Mineral oil; HC: Hydrogen cyanamide; B: Bluprins[®]; CV: coefficient of variation; ^{ns}: not significant at 5% probability

For 'Fuji Suprema' apple trees and fruit production per tree, the treatments Bluprins[®] 3% + Ca(NO₃)₂ 3% and Bluprins[®] 3% + Ca(NO₃)₂ 3%+ NH₄(NO₃) 3% resulted in lower fruit production per tree than that of other treatments and did not differ from the control treatment in the 2013/2014 growing season. In the 2014/2015 growing season, except for the treatment Bluprins[®] 3% + Ca(NO₃)₂ 3%+ NH₄(NO₃) 3%, the treatments resulted in lower production of fruit per tree in compared to the control treatment. In 2015/2016, the harvest was not evaluated and in 2016/2017, the treatments did not differ from each other (Table 10).

According to Botelho et al. [33], the response of bud break promoters on fruit production is dependent on the climatic conditions during the fruit development period. Once the climatic conditions were adequate during this period in the growing seasons studied, there was no marked difference in the fruit production per tree by the use of bud break promoters, for both cultivars.

The average fruit weight did not showed significant differences between treatments in the 2013/2014 and 2016/2017 growing seasons for 'Maxi gala', and for 'Fuji Suprema' cultivar and insignificant between treatments in 2013/2014, 2014/2015 and 2015/2016. For 'Maxi Gala' apple trees in the 2014/2015 growing season, the treatments Bluprins[®] 5% + Ca(NO₃)₂ 3%, Bluprins[®] 3% + Ca(NO₃)₂ 3%+ NH₄(NO₃) 3% and Bluprins[®] 5% + Ca(NO₃)₂ 4%+ NH₄(NO₃) 4% did not differ from the control treatment and resulted in lower values than the other treatments (Tables 9 and 10).

It was observed that there were some differences between 'Maxi Gala' and 'Fuji Suprema' cultivars, such as the flowering period, lower in 'Maxi Gala', and the percentage of buds sprouted, higher in 'Fuji Suprema'. However, both cultivars presented similar kind of behaviour in response to climatic factors such as the anticipated beginning of bud break and flowering in the growing seasons with colder winter, and higher bud break by the application of bud break

Table 9. Fruit production per plant (FPP, kg) and average fruit weight (AFW, g) of 'Maxi Gala' apple trees under the influence of compounds for bud break during tree growing seasons. Caçador, SC, 2018

Treatments	2013/2014		2014/2015		2016/2017	
	FPP (kg)	AFW (g)	FPP (kg)	AFW (g)	FPP (kg)	AFW (g)
1. Control	9.0 b	145.1 ^{ns}	8.0 ^{ns}	120.9 b	4.7 b	132.8 ^{ns}
2. MO 3,5%+HC 0,35%	3.9 b	137.4	13.1	138.0 a	8.4 a	139.7
3. B 3% + Ca(NO ₃) ₂ 3%	9.4 b	165.3	10.1	133.0 a	5.0 b	154.3
4. B 5% + Ca(NO ₃) ₂ 3%	13.9 a	141.3	13.6	118.6 b	4.9 b	148.3
5. B 3% + Ca(NO ₃) ₂ 5%	16.1 a	142.6	11.0	130.3 a	3.7 b	149.4
6. B 5% + Ca(NO ₃) ₂ 5%	9.6 b	149.6	14.6	130.1 a	6.8 b	151.8
7. B 3%+Ca(NO ₃) ₂ 3%+NH ₄ (NO ₃) 3%	19.9 a	140.6	15.2	121.4 b	10.5 a	150.1
8. B 5%+Ca(NO ₃) ₂ 4%+NH ₄ (NO ₃) 4%	16.6 a	140.0	12.8	119.2 b	6.0 b	140.9
CV (%)	33.2	11.2	39.8	8.1	52.1	12.4

MO: Mineral oil; HC: Hydrogen cyanamide; B: Bluprins[®]; CV: coefficient of variation. Means followed by same letter do not differ by Scott-Knott test at 5% probability. ns: not significant

Table 10. Fruit production per plant (FPP, kg) and average fruit weight (AFW, g) of 'Maxi Gala' apple trees under the influence of compounds for bud break during tree growing seasons. Caçador, SC, 2018

Treatments	2013/2014		2014/2015		2016/2017*	
	FPP (kg)	AFW (g)	FPP (kg)	AFW (g)	FPP (kg)	AFW (g)
1. Control	25.9 b	123.6 ^{ns}	35.1 a	129.2 ^{ns}	19.6 ^{ns}	101.7 ^{ns}
2. MO 3,5%+HC 0,35%	35.6 a	113.9	19.7 b	120.5	11.8	118.2
3. B 3% + Ca(NO ₃) ₂ 3%	23.4 b	116.7	15.4 b	138.0	14.6	102.2
4. B 5% + Ca(NO ₃) ₂ 3%	34.5 a	120.3	23.0 b	124.8	15.4	111.1
5. B 3% + Ca(NO ₃) ₂ 5%	40.6 a	118.7	17.8 b	126.6	14.4	104.6
6. B 5% + Ca(NO ₃) ₂ 5%	42.7 a	113.6	19.6 b	125.5	17.9	102.5
7. B 3%+Ca(NO ₃) ₂ 3%+NH ₄ (NO ₃) 3%	14.2 b	125.9	37.3 a	122.7	11.8	114.8
8. B 5%+Ca(NO ₃) ₂ 4%+NH ₄ (NO ₃) 4%	38.0 a	111.3	13.3 b	122.1	15.2	112.0
CV (%)	31.9	118.0	22.7	126.2	33.1	12.2

MO: Mineral oil; HC: Hydrogen cyanamide; B: Bluprins[®]; CV: coefficient of variation. Means followed by same letter do not differ by Scott-Knott test at 5% probability. ns: not significant

promoters, especially mineral oil + hydrogen cyanamide.

4. CONCLUSION

Bluprins[®] in combination with calcium nitrate and ammonium nitrate proved effective in inducing bud break of 'Maxi Gala' and 'Fuji Suprema' apple tree cultivars under mild winter conditions, however, this is less effective than mineral oil + hydrogen cyanamide. Bluprins[®] in combination with calcium nitrate and ammonium nitrate anticipates the bud break and flowering period and reduces the flowering period for 'Maxi Gala' and 'Fuji Suprema' apple tree cultivars, in compared to plants without application of bud break promoters. Bluprins[®] does not affect the fruit set considerably and fruit production of 'Maxi Gala' and 'Fuji Suprema' apple tree cultivars.

COMPETING INTERESTS

Authors have declared competing interest for the company name "Bluprins[®]".

REFERENCES

- Food and Agriculture Organization of the United Nations Statistical Databases. Crops. (Accessed 15 August 2018)
Available: <<http://www.fao.org/faostat/en/#data/QC>>
- Associação Brasileira de Produtores de Maçã. Anuário Brasileiro da Maçã; 2016. (Accessed 15 August 2018)
Available: <<http://www.abpm.org.br/anuario-s-da-maca>>
- Campoy JA, Ruiz D, Egea J. Dormancy in temperate fruit trees in a global warming

- context: A review. *Sci. Hortic.* 2011;130: 357-372.
DOI: 10.1016/j.scienta.2011.07.011.
4. Atkinson CJ, Brennan RM, Jones HG. Declining chilling and its impact on temperate perennial crops. *Environ Exp Bot.* 2013;91:48-62.
DOI: 10.1016/j.envexpbot.2013.02.004
 5. Cornille A, Gladieux P, Smulders MJM, Roldán-Ruiz I, Laurens F, Le Cam B, et al. New insight into the history of domesticated apple: Secondary contribution of the European wild apple to the genome of cultivated varieties. *PLoS Genet.* 2012;8(5):e1002703.
DOI: 10.1371/journal.pgen.1002703
 6. Petri JL, Leite GB, Couto M, Francescato P. Avanços na cultura da macieira no Brasil. *Rev Bras Frutic.* 2011;33:48-56. Portuguese.
DOI:10.1590/S0100-29452011000500007
 7. Petri JL, Leite GB. Consequences of insufficient winter chilling on apple tree bud-break. *Acta Hortic.* 2004;662(1):53-60.
DOI: 10.17660/ActaHortic.2004.662.4
 8. Anzanello R, Fialho BF, Santos HP, Bergamaschi H, Marodin GAB. Bud dormancy in apple trees after thermal fluctuation. *Pesqui Agropecu Bras.* 2014;49(6):457-464.
DOI:10.1590/S0100-204X2014000600007
 9. Hawerth FJ, Petri JL, Leite GB. Cianamida hidrogenada, óleos mineral e vegetal na brotação de gemas e produção de macieiras 'Royal Gala'. *Semin Cienc Agrar.* 2010;31(1):1145-1154. Portuguese.
DOI:10.5433/1679-0359.2010v31n4Sup1p1145
 10. Pasa MS, Felippeto J, Nava G, Silva CP, Brighenti AF, Ciotta MN. Performance of 'Fuji Suprema' apple trees treated with budbreak promoters, in São Joaquim-SC. *Rev Bras Frutic.* 2018;40(1).
DOI: 10.1590/0100-29452018325
 11. Rufato L, Kretzschmar AA, Brighenti AF, Macedo TA, Mendes M, Silva LC. Bud break in different cultivars of apple trees in two regions of Santa Catarina State, Brazil. *Acta Hortic.* 2010;884:643-646.
DOI: 10.17660/ActaHortic.2010.884.85
 12. Petri JL, Leite GB, Couto M, Gabardo GC, Hawerth FJ. Chemical induction of budbreak: new generation products to replace hydrogen cyanamide. *Acta Hortic.* 2014;1042:159-166.
DOI: 10.17660/ActaHortic.2014.1042.19
 13. Campoy JA, Ruiz D, Egea, J. Effects of shading and thidiazuron + oil treatment on dormancy breaking, blooming and fruit set in apricot in a warm-winter climate. *Sci. Hortic.* 2010;125(3):203-210.
DOI: 10.1016/j.scienta.2010.03.029
 14. Erez A. Bud dormancy, phenomenon, problems and solutions in the tropics and subtropics. In: Erez A, editor. *Temperate Fruit Crops in Warm Climates.* Dordrecht: Springer; 2000.
(Accessed 12 march 2014)
Available:https://link.springer.com/chapter/10.1007/978-94-017-3215-4_2
 15. Petri JL, Couto M, Leite GB, Gabardo GC, Francescato P. Tratamentos alternativos para indução da brotação e floração na macieira. In: *Anais do XXII Congresso Brasileiro de Fruticultura.* Bento Gonçalves: Embrapa; 2012.
 16. Mohamed AKA. The effect of chilling, defoliation and hydrogen cyanamide on dormancy release, budbreak and fruiting of Anna apple cultivar. *Sci Hortic.* 2008;118(1):25-32.
DOI: 10.1016/j.scienta.2008.05.015
 17. Hawerth FJ, Petri JL, Herter FG, Leite G B, Leonetti JF, Marafon AC, et al. Fenologia, brotação de gemas e produção de frutos de macieira em resposta à aplicação de cianamida hidrogenada e óleo mineral. *Bragantia.* 2009;68(4):961-971. Portuguese.
DOI:10.1590/S0006-87052009000400016
 18. Hawerth FJ, Petri JL, Leite GB, Herter FG. Brotação de gemas em macieiras 'Imperial Gala' e 'Fuji Suprema' pelo uso de Erger e Nitrato de Cálcio. *Rev Bras Frutic.* 2010;32(2):343-350. Portuguese.
DOI:10.1590/S0100-29452010005000074
 19. Ziosi V, Giovannetti G, Vitali F, Di Nardo A, Costa G, Laghezza L. Bluprins® induces advanced bud release from dormancy in sweet cherry. *Acta Hortic.* 2013;1009:71-77.
DOI: 10.17660/ActaHortic.2013.1009.7
 20. Ziosi V, Di Nardo A, Fontana A, Vitali F, Costa G. Effect of Bluprins® Application on Bud Release from Dormancy in Kiwifruit, Cherry, and Table Grape. In: Anderson JV, editor. *Advances in Plant Dormancy.* Switzerland: Springer International Publishing; 2015.
 21. Sanhueza RMV, Protas JFS, Freire JM. Manejo da Macieira no Sistema de Produção Integrada de Frutas. Bento Gonçalves: Embrapa Uva e Vinho; 2006.

22. Petri JL, Leite GB, Putti GL. Apple tree bud break promoters in mild winter conditions. *Acta Hortic.* 2008;774:291-296. DOI: 10.17660/ActaHortic.2008.774.39
23. Shaltout AD, Unrath CR. Rest completion prediction model for 'Starkrimson Delicious' apples. *J Am Soc Hortic Sci.* 1983;108(6): 957-961.
24. Ferreira DF. *Sisvar*, version 5.6. Universidade Federal de Lavras, Lavras; 2010.
25. Kozma P, Nyéki J, Soltész M, Szabó Z. *Floral Biology, Pollination and Fertilization Zone Fruit Species and Grape.* Budapest: Akademiai Kiado; 2003.
26. Petri JL. Práticas de manejo da quebra de dormência em pomáceas. In: *Trabajos presentados in Seminario de Actualización Técnica Frutales de Pepita, Programa de Investigación em Producción Frutícola, Las Brujas: INIA; 2014. Uruguay.*
27. Hawerth FJ, Petri JL, Herter FG, Leite GB, Leonetti JF, Marafon AC, et al. Fenologia, brotação de gemas e produção de frutos de macieira em resposta à aplicação de cianamida hidrogenada e óleo mineral. *Bragantia.* 2009;68:961-971. Portuguese. DOI:10.1590/S0006-87052009000400016
28. Leite GB, Bonhomme M, Putti GL, Petel G, Petri JL, Rageau R. Physiological and biochemical evolution of peach leaf buds during dormancy course under two contrasted temperature patterns. *Int J Hortic Sci.* 2006;12(4):15-19.
29. Marchi T, Oliari CR, Maia AJ, Sato AJ, Botelho RV. Indução da brotação de gemas de macieiras com aplicação de óleos vegetais e mineral. *Ver. Ciênc. Agron.* 2017;48(3):501-512. Portuguese. DOI: 10.5935/1806-6690.20170059
30. Cruz Júnior AO, Ayub RA. Quebra de dormência de gemas de macieira cv. Eva tratadas com cianamida hidrogenada. *Rev. Bras. Frutic.* 2002;24(2):576-578. Portuguese. DOI: 10.1590/S0100-29452002000200061
31. El-Agamy SZ, Mohamed AKA, Mostafa FMA, Abdallah AY. Effect of GA₃, Hydrogen Cyanamid and Decapitation on Budbreak and Flowering of Two Apple Cultivars under the Warm Climate of Southern Egypt. *Acta Hortic.* 2001;565: 109-114. DOI: 10.17660/ActaHortic.2001.565.17
32. Ramirez F, Davenport TL. Apple pollination: a review. *Sci. Hortic.* 2013; 162(23):188-203. DOI: 10.1016/j.scienta.2013.08.007
33. Botelho RV, Pires EJP, Terra MM. Brotação e produtividade de videiras da cultivar Centennial Seedless (*Vitis vinifera* L.) tratadas com cianamida hidrogenada na região noroeste do Estado de São Paulo. *Rev Bras Frutic.* 2002;24(3):611-614. Portuguese. DOI: 10.1590/S0100-29452002000300007

© 2018 Fenili et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/26381>