



Geological Dynamics and Stability in Lansdowne, Garhwal Himalayas: A Litho- structural Perspective

**Kartik Patel ^{a,b,*}, Mridul Anand ^{a,c}
and Shaurya Vikram Singh ^{a,c}**

^a Department of Geology, Babasaheb Bhimrao Ambedkar University, Lucknow-226025, Uttar Pradesh, India.

^b Department of Earth and Environmental Studies, National Institute of Technology, Durgapur-713209, West Bengal, India.

^c IHEG, ONGC Center of Advanced Studies, Lucknow University, Lucknow-226007, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/120277>

Original Research Article

Received: 20/05/2024
Accepted: 24/07/2024
Published: 29/07/2024

ABSTRACT

This study examines the geological features of the Lansdowne town in the Lesser Himalayas of Uttarakhand, India, where the rocks ranging from Precambrian to Recent are present. The Lansdowne region showcases intricate geological features, encompassing various stages of deformation, and notable compositional changes from augen gneiss to granite as one approaches the town's core. Detailed field analyses of the rock formations and geological structures suggest significant tectonic movements which are marked by the presence of fault lines, joint sets, and gouge

*Corresponding author: Email: kartikpatel171101@gmail.com;

Cite as: Patel, Kartik, Mridul Anand, and Shaurya Vikram Singh. 2024. "Geological Dynamics and Stability in Lansdowne, Garhwal Himalayas: A Litho- Structural Perspective". *Asian Journal of Geological Research* 7 (2):175-83. <https://journalajoger.com/index.php/AJOGER/article/view/163>.

zones. The lithological variation from the highly distorted augen gneiss to a more uniform granitic core helps to understand the different tectonic activities and metamorphic conditions in the area. The central part of Lansdowne, which is characterized by uniform granite, shows less intense deformation, suggesting a more stable environment for development. This study highlights the importance of geological stability in urban planning within seismically active regions.

Keywords: Lesser Himalaya; lansdowne; granite; gneiss, fault.

ABBREVIATIONS

MFT : The Main Frontal Thrust
HFT : Himalayan Frontal Fault
MBT : Main Boundary Thrust
MCT : Main Central Thrust
STDS : South Tibetan Detachment System
GPS : Global Positioning System
GIS : Geographic Information System

1. INTRODUCTION

The Himalayas, located on the southern fringe of the Tibetan Plateau, are the youngest mountain range and stretches from west to east [1]. The mountain range is arc-shaped, convex southwards with syntaxial bends at the western and eastern ends [2]. They are formed as a result of the collision between the Indian and Eurasian plates [3]. Due to ongoing continental collision, the Himalayas stand as the Earth's highest mountain range undergoing active uplift. Himalaya stretches about 2500km in length from Nanga Parbat (8138m) in the northwest to Namcha Barwa (7756m) in the northeast [4]. The width varies from 350 km in the west (Kashmir) to 150 km in the east (Arunachal Pradesh) [5]. During the convergence, the large-scale crustal shortening was accommodated by south directed deformation related tectonic transport of thrust sheets all along the length of the Himalayas [6,7]. As a result of the subduction of sediments due to the collision between the plates, this mountain range shows an agreement among various dating methods indicating that the older segments originated during the late Cretaceous period (82-113 million years ago), while the newer areas were formed in the early Eocene epoch (40-60 million years ago).

The Himalayan Mountain range is subdivided into four principal tectonic zones, from south to north: the Sub- Himalaya, the Lesser Himalaya, the Higher Himalayan Crystallines, and the Tethyan Himalaya. The generalized geological succession of the Himalayas is shown in Table 1.

- **Sub-Himalaya (Siwalik range or Outer Himalaya):** It has an average altitude of 500m above sea level. The Main Frontal Thrust (MFT) or Himalayan Frontal Fault (HFT), separates the outer Himalaya from the Ganga plains [8]. It consists predominantly of Tertiary and Quaternary sediments [2]. It is bounded to the north by a northward dipping Main Boundary Thrust (MBT), separating it from the overlying Lesser Himalaya [8].
- **Lesser Himalaya (Lower or Middle Himalaya):** It is higher than 2500 m, however, in the Kashmir Valley, the Pir Panjal Range rises to heights greater than 3500 m [2]. It consists of Precambrian and Cambrian sequences of the Damtha, Tejam, Jaunsar-Garhwal, and Mussoorie Groups in the Garhwal region of the western Himalaya [9]. The ranges of the lower Himalaya are i) Pir Panjal range, ii) Dhaola Dhar range, iii) Mussoorie and Nag Tibba range, and iv) Mahabharat Lekh range [10].
- **Higher Himalaya (Greater or Inner or Central Crystalline Himalaya):** It is separated from the Lesser Himalaya along the Main Central Thrust (MCT). It has an average elevation of 6000 m above mean sea level [8]. The Higher Himalayan Crystallines in the western Himalaya is represented by the Chail, Jutogh, and Vaikrita groups of rocks [2]. It is incredibly rugged with perennial snow and glacier-capped peaks [11].
- **Trans Himalaya (Tethys or Tethyan Himalaya):** It is rugged terrain with an average elevation of 3000 m above mean sea level [8]. The Tethyan Himalayan sequence is separated from the Higher Himalayan Crystalline sequence by South Tibetan Detachment System (STDS) [12]. The high ranges of the Higher Himalayan Crystalline act as a barrier for the wet Monsoon cloud to go further north, the Tethyan Himalayan Range has developed

into a cold and dry desert with sparse vegetation and naked rocks [2].

Fig. 1 shows the divisions of the Himalayas and Fig. 2 shows the cross-section of the Himalayas.

1.1 Geological Setting

The Garhwal Himalayan forms a part of the Lesser Himalayas, bounded by the Main Boundary Thrust (MBT) in the south and the Main Central Thrust (MCT) in the north [13]. Lansdowne town is situated in the Pauri Garhwal district of Uttarakhand state. The Lansdowne granitic gneisses form a part of the Amri tectonic unit of Garhwal nappe [14]. The area is characterized by a tectonic contact between the Lansdowne granite gneiss and the underlying Lansdowne metamorphites [15]. These metamorphites have undergone polyphase progressive regional metamorphism,

as evidenced by three distinct deformational and folding events [16]. A unique assemblage of minerals including chloritoid, staurolite, kyanite, andalusite, and quartz is present within the region. The granitic rocks themselves have been metamorphosed into augen gneiss, mylonites, and phyllonites, indicating significant deformation [15]. Additionally, geochemical alteration of the granitic rocks is evident by processes like muscovitization, tourmalinization, and greisenization. These textural and geochemical features within the rocks can be directly linked to the deformational and folding events that have shaped the Lansdowne area. The granitic gneisses are well foliated. In the northwest they extend up to Bhayansu on the steep slopes of Lansdowne hill. They occupy the core of the Lansdowne synform. Litho-tectonic succession in the Lansdowne area is shown in Fig 3 and the geological map of the Lansdowne area is shown in Fig. 4.

Table 1. Generalized geological succession of the Himalayas

Indus Tsangpo Suture Zone (ITSZ)	
Late Precambrian- Cenozoic	Trans Himalaya
South Tibetan Detachment Zone (STDZ) or Tethys Thrust	
Precambrian	Greater Himalaya
Main Central Thrust (MCT)	
Precambrian	Lesser Himalaya
Main Boundary Thrust (MBT)	
Tertiary	Siwalik Himalaya
Main Frontal Thrust (MFT)/ Himalayan Frontal Thrust (HFT)	
Quaternary	Indo-Gangetic Plain




Fig. 1. The divisions of Himalayas




Fig. 2. Cross section of Himalayas




Fig. 3. Litho-tectonic succession in the Lansdowne area [17]




Fig. 4. Geological map of Lansdowne area (after Fuchs and Sinha, 1978)

2. MATERIALS AND METHODS

The research involved conducting thorough fieldwork to collect data on geological structures, compositional variations, and lithological changes in the Lansdowne area. Initially, potential sites were identified using topographic and geological maps, with a focus on accessibility and geological significance. At every location, a Global Positioning System (GPS) was used to accurately determine the latitude and longitude, with the coordinates recorded in a notebook alongside the site identification numbers. The foliation trends of rock outcrops were assessed using a Brunton compass, and several measurements were taken to ensure precision. A 10x hand lens was employed to analyze the mineral compositions, pinpointing significant minerals and their interrelationships. Geological maps were consulted to identify representative outcrops, and the locations of these outcrops were marked using GPS coordinates. Rock samples were then collected from these specific locations. Detailed notes were made on the changes in rock types, and photographs were captured of significant geological features. Finally, the gathered data were incorporated into a Geographic Information System (GIS) for spatial analysis, enabling precise mapping of geological features.

3. RESULTS AND DISCUSSION

Through field investigations and petrographic analyses, we have discovered notable differences in the structural and lithological characteristics of the Lansdowne region within the Garhwal Himalayas. The key findings are as follows:

- Lithological Variation:** Towards the center of Lansdowne town particularly noteworthy, there is a noticeable shift in the type of rock from augen gneiss to granite. This transitional area is characterized by a gradual change in the rock's structure, with the foliated layers of gneiss gradually fading away, and the composition shifting towards a more granitic nature. Augen gneiss, characterized by its prominent large feldspar crystals, suggests a significant level of metamorphism and deformation. The shift to granite may reflect a decrease in the intensity of deformation and metamorphism towards the core of the area, or it could indicate different stages of intrusion and cooling of magmatic bodies. The granite is composed of quartz, alkali feldspar, plagioclase, muscovite, biotite, and tourmaline. Fig. 5 (a-b) shows the augen gneiss and the granite found in the study area and Fig. 6 shows the traversing map.

- **Faults:** A fault is a fracture with appreciable displacement parallel to the fracture plane or fault surface. A fracture occurs when a rock breaks in response to stress and divides it into two or more pieces [18]. Local quartz veins show small-scale faulting. A reverse fault is observed in the study area in which the hanging wall block moves up relative to the footwall block (Fig. 6). The presence of faults that the Himalayas have experienced multiple periods of deformation, indicating a dynamic tectonic environment.
- **Gouges:** A fault gouge is a fine-grained substance near the surface, which is essentially the rock powder formed during faulting [19]. Fragments of finely crushed rock and clay minerals were discovered along the fault planes, indicating the presence of gouge material (Fig. 7). The presence of gouges along the fault planes is indicative of significant brittle deformation and substantial movement. It also suggests the occurrence of fluid-assisted alteration processes during faulting.
- **Joint Sets:** Joints are the fractures or cracks along which no displacement occurs [18]. A joint set is a group of parallel fractures with a common origin [20]. Three different sets of joints were identified (Fig. 8). The orientation of the joint sets suggests that the area has been subjected to varying stress regimes, which is consistent with the ongoing tectonic activity in the region.




Fig. 5. (a) Augen gneiss, (b) Granite




Fig. 6. Traversing Map



Fig. 7. Reverse fault



Fig. 8. Fault gouge



Fig. 9. Three sets of joints

4. CONCLUSION

This geological study conducted in the Lansdowne region of the Garhwal Himalayas has uncovered the intricate geological complexities, encompassing the various phases of the deformation, faulting, and compositional changes from the augen gneiss to granite. The area has a tectonic contact between the Lansdowne granite gneiss and the underlying Lansdowne metamorphites. The presence of the faults, joint sets, and gouge zones suggest significant tectonic activity, while the transition from highly deformed augen gneiss to the more uniform granite towards the core of Lansdowne indicates different tectonic and metamorphic conditions in the region. The research indicates that the central part of the Lansdowne area which is known for its uniform granite composition, has experienced less intense deformation and metamorphism compared to the surrounding regions dominated by augen gneiss. The relative stability and lower deformation intensity in the granitic core suggest a potentially safer and more stable environment for the establishment of settlements.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. De Jong W. Chapter 1 - An overview of the Himalaya forests and their drivers of change," in Climate Change in the Himalayas, A. Kumar, W. D. Jong, M. Kumar, and R. Pandey, Eds.: Academic Press, 2023;1-11.
2. Ahmad TJG. Geology of the Himalayan mountain range, with special reference to the western Himalaya. 2011;63; 142-147.
3. Molnar P, Tapponnier PJS. Cenozoic tectonics of Asia: Effects of a continental collision: Features of recent continental tectonics in Asia can be interpreted as results of the India-Eurasia collision. 1975;189(4201):419-426.
4. Roy AB, Purohit R. Chapter 17 - The Himalayas: Evolution Through Collision, in Indian Shield, A. Roy B, Purohit R. Eds.: Elsevier, 2018;311-327.
5. Karan P. Geographic Regions of the Himalayas," 05/24 2012.
6. KJPT, OT, RS, OLS, Valdiya A, Mathematical P. Sciences, Tectonics and evolution of the central sector of the Himalaya. 1988;326(1589):151-175.
7. Kapp P, PG, JA, JO, DeCelles S. Mesozoic–cenozoic geological evolution of the Himalayan- Tibetan orogen and working tectonic hypotheses. 2019;319 (3):159-254.
8. VJLHM, ITH, KHH, Thakur DO. Regional geology and geological evolution of the Himalaya. 2001;53055.
9. Bhargava ON. The precambrian sequences in the western Himalayas," Geol. Surv. India Spec. Publ. 2000;55:69-84.01/01 2000.
10. Valdiya KS, JNC, FAS. Research, dynamic Himalaya. Jawaharlal Nehru Centre for Advanced Scientific Research; 1998.
11. Sharma UC, Datta M, Sharma V. Soils in the Hindu Kush Himalayas: management for agricultural land use. Springer; 2022.
12. Carosi R, Lombardo B, Molli G, Musumeci G, Pertusati PC. The South Tibetan detachment system in the Rongbuk valley, Everest region. Deformation features and geological implications. Journal of Asian Earth Sciences. 1998;16(2):299-311. 1998/04/01/ 1998.
13. Rawat A, Banerjee S, Sundriyal YJG, Geomorphological and statistical assessment of tilt-block tectonics in the garhwal synform: Implications for the active tectonics. Garhwal Lesser Himalaya, India. 2021;11(8):345.
14. Shah A, PJG, SOI, Patel. granitic gneisses of Lansdowne (District Pauri-Garhwal, UP). 1978;19(8):368-372.
15. LJ, GS, OI. Gupta. A contribution to the Geology of Lansdowne area, Garhwal Himalayas, India. 1976;17(4):449-460,
16. Chakrabarti BK. Precambrian geotectonics in the Himalaya: Sans cenozoic hangover. Elsevier; 2023.
17. Maity S, Rawat A, Banerjee S, Srivastava H. Structural architecture of the Lansdowne thrust and Garhwal Nappe in Lansdowne area of western Himalaya; 2018.

18. Gangwar I, Chapter: 12 Geology of Gwalior and Adjacent Areas, Madhya Pradesh (India) Isha Gangwar Department of Geology, Babasaheb Bhimrao Ambedkar University, Lucknow-226025, Uttar Pradesh, India.
19. Higgins MW. Cataclastic rocks. US Government Printing Office; 1971.
20. Liu E, Martinez A. 1 - Introduction, in Seismic Fracture Characterization, E. Liu and A. Martinez, Eds. Oxford: EAGE, 2012;13-27.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/120277>