



# ***In vitro* Investigation of Phytoconstituents, GC-MS, TLC, Antioxidant activity, Total Phenolic & Flavonoid Contents from *Aegle marmelos L* (Bael) Leaves Extract**

**Ramesh Kumar <sup>a+++</sup>, Smrati Sharma <sup>a</sup>, Sidhanshu Kumar <sup>a</sup>,  
Dinesh Kumar <sup>b</sup>, Rekha Lagarkha <sup>c</sup>, Sanjay Kumar <sup>d</sup>  
and Maneesha Pandey <sup>e</sup>**

<sup>a</sup> Department of Biochemistry, Bundelkhand University, Jhansi (UP)- 284128, India.

<sup>b</sup> Department of Chemistry, Bipin Bihari College, Jhansi (UP), India.

<sup>c</sup> Department of Chemistry, Bundelkhand University, Jhansi (UP)- 284128, India.

<sup>d</sup> Department of Microbiology, Bundelkhand University, Jhansi (UP)- 284128, India.

<sup>e</sup> Biochemistry Discipline, School of Sciences, Indira Gandhi National Open University,  
New Delhi 110068, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/ejmp/2024/v35i61218>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124581>

**Original Research Article**

**Received: 14/08/2024**  
**Accepted: 17/10/2024**  
**Published: 24/10/2024**

<sup>++</sup>Assoc. Prof. & Former-Head;

\*Corresponding author: E-mail: [drramesh.kumar34@gmail.com](mailto:drramesh.kumar34@gmail.com);

**Cite as:** Kumar, Ramesh, Smrati Sharma, Sidhanshu Kumar, Dinesh Kumar, Rekha Lagarkha, Sanjay Kumar, and Maneesha Pandey. 2024. "In Vitro Investigation of Phytoconstituents, GC-MS, TLC, Antioxidant Activity, Total Phenolic & Flavonoid Contents from *Aegle Marmelos L* (Bael) Leaves Extract". *European Journal of Medicinal Plants* 35 (6):187-99. <https://doi.org/10.9734/ejmp/2024/v35i61218>.

## ABSTRACT

The Rutaceae family includes *Aegle marmelos* L., a moderately sized aromatic tree that is also known as Bael, Bengal quince, golden apple, Japanese bitter orange, stone apple, or wood apple. It is mostly found in tropical or subtropical regions. Fever, seminal weakness, nausea, vomiting, swellings, diarrhea, dyspepsia, and intermittent fever are the most common ailments that bael leaves are used to treat. Because of their high phenolic and flavonoid content, leaves have anti-allergic, anti-inflammatory, anti-microbial, anti-thrombotic, cardioprotective, and vasodilatory properties. Therefore, the aim of the present study was to determine the phytochemical components, antioxidant capacity, GC-MS analysis, total phenolic & flavonoid contents in the methanolic extracts of *A. marmelos* leaves. Various secondary metabolites were observed in the aqueous and methanolic *A. marmelos* leaf extracts. TLC analysis of methanolic extracts revealed five distinct spots having different R<sub>f</sub> values. Total phenolic content was found to be 33 ± 7.62 mg GAE/g, flavonoid content was 307.8 ± 130.12 mg QE/g. The highest antioxidant activity of methanolic extract of *A. marmelos* leaves was obtained at 1000 µg/ml conc. and it was dose dependent. Many biologically active compounds were present as analysed by GC-MS. The highest peak area of Perylo[1,12-def]-1,3-dioxepin-5,11-dione, 6,12-dihydroxy-8,9-bis(2-hydroxypropyl)-7,10-dimethoxy-, stereoisomer (3.502%) was observed.

**Keywords:** *A. marmelos*; ascorbic acid; antioxidant activity; GC-MS; total phenolic content; total flavonoid content.

## 1. INTRODUCTION

*Aegle marmelos* L., a perennial tree belonging to the Rutaceae family, is widely distributed in sub-Himalayan region, central and southern India. It is commonly called as Vilvam in Tamil, Bilva in Sanskrit, and Bael in Hindi. Its leaves are utilized in spiritual practices to honor Lord Shiva. It is widespread to India and is utilized in folk medicines. Plants are excellent sources for the investigation of pharmaceutical compounds and drugs. Natural products and their analogues can serve as intermediates in the manufacture of beneficial pharmaceuticals, and they may also have therapeutic value for humans or live stock species [1].

In developing nations, traditional medical practices mostly the use of medicinal plants remain essential for providing basic healthcare needs. The therapeutic usefulness of these plants is because of the presence of active chemical components (phytoconstituents) that leads to physiological action on the human body. Alkaloids, tannin, flavonoids, and phenolic compounds are the most significant of these bioactive components of plants [2], along with carotenoids, ascorbic acid, tocopherol, and polyphenols [3-5].

“The use of synthetic antioxidants leads many adverse effects while natural antioxidants are safe with no side effects” [6]. “Antioxidant substances prevent damages caused by free

radicals. Apoptosis, gene expression, and cell proliferation are just a few of the regulatory mechanisms that are impacted by free radicals produced during aerobic metabolism. Excessive generation of free radicals may disrupt the antioxidant defence systems, damaging vital biomolecules (such as membrane lipids, cell proteins, carbohydrates, DNA, and enzymes). Moreover, oxidative stress causes the occurrence of cytotoxic compounds and modifies the oxidant-antioxidant balance (redox homeostasis) that characterizes normal cell functioning” [7-9]. Pathologies caused by oxidative stress include cancer [10,11], cardiovascular disease [12], neural disorders [13], Alzheimer’s disease [14], mild cognitive impairment [15], Parkinson’s disease [16], alcohol-induced liver disease [17], ulcerative colitis [18], atherosclerosis [19], and aging [20]. Polyphenols, flavonoids, vitamin C and Vitamin E are phytochemicals in foods that have been known as a natural antioxidant and shows antioxidant activity. Polyphenols and flavonoids act as reactive oxygen species (ROS) scavenger and have chemopreventive activity [21,22]. Ageing, atherosclerosis, and cancer are only a few of the pathological processes that are impacted by ROS [23]. Thus, scavenging ROS can minimize chronic illness. It has been reported that taking foods high in polyphenols and flavonoids contents may reduces the inflammation, oxidative stress, tumors, and coronary heart disease [22,24,25]. Antioxidant functions in reducing DNA damage, lowering lipid

peroxidation, supporting the immune system and inhibit the malignant transformation of cells [26]. Plants that contain polyphenolic chemicals such as flavonoids have antioxidant potential [27]. According to prior studies, the biological actions of many compounds are related to their antioxidant activity. In this present investigation, phytochemical screening of aqueous and methanolic extracts of leaves of *A. marmelos* was evaluated and biologically important compounds were identified by GC-MS analysis. Thin Layer Chromatography, total phenolic & flavonoids contents, and antioxidant activity of *A. marmelos* leaves was determined.

## 2. MATERIALS AND METHODS

### 2.1 Plant Materials

Leaves sample of *A. marmelos* was collected from the Bundelkhand University campus, Jhansi and was identified by Dr. J. C. Arya Research Officer, (Botany) from the Central Ayurveda Research Institute under the Central Council for Research in Ayurvedic Sciences, Ministry of AYUSH, Gwalior Road, Jhansi (Uttar Pradesh). The leaves were thoroughly washed with tap water followed by de-ionized water. After washing, it was kept in the dark for drying at room temperature and finally crushed with the help of mixer & grinder. The extraction was performed with the stored plant materials.

### 2.2 Extraction Procedure

Leaves sample was extracted by two methods –

- Aqueous extraction and
- Methanolic extraction

**Aqueous Extract:** In the water bath, different amounts of powdered sample i.e. 5 and 10 gms were extracted for one hour at 90°C with 100 ml of de-ionized water. After 1 hour, the extract was filtered and stored at 4°C for the further tests.

**Methanolic Extract:** The powdered leaves were percolated using 80% of methanol in the Soxhlet apparatus at 60-65°C. This extract was evaporated to dryness in a water bath at 40°C temperature and stored in air tight bottles.

### 2.3 Phytochemical Analysis

The presence or absence of secondary metabolites was carried out as described previously [28-30]. The alkaloids, carbohydrates,

reducing sugars, flavonoids, glycosides, tannin, saponin, protein, triterpenoids & steroids were determined in the aqueous and methanolic extracts of leaves of *A. marmelos*. More detail procedures are mentioned elsewhere [28].

### 2.4 Total Antioxidant Capacity (TAC) by Phosphomolybdenum Assay

“The total antioxidant activity was measured using the phosphomolybdenum test [31]. Two milliliters of the reagent solution were mixed with different conc. of the test sample (1000-31.25 µg/ml). The reaction mixture was incubated for 90 minutes at 95°C in an oven-cum-incubator. For the blank tube, the extract was replaced with 80% methanol in the same ratio. Ascorbic acid was used as a reference control. The samples were cooled at room temperature and the absorbance taken at 695 nm wavelength using a multiplate reader” [32].

### 2.5 Thin layer chromatography

“Methanolic extract of *A. marmelos* leaves was tested on TLC plates coated with 0.2 mm thick silica gel-G. The solvent mixture that we employed in this instance was butanol, acetic acid, and water at a 2:1:1 v/v ratio, as previously mentioned. Spots move with the solvent mixture on the silica-coated plates due to capillary action, detecting the methanolic extract over 4 mm from the plate base. A fully developed silica coated plate was heated for 20-25 minutes after being allowed to air dry. To observe the bands, 0.2% freshly prepared ninhydrin solution was sprayed across the plate” [32].

These spots were expressed by its retention factor (Rf).

$$R f = \text{Distance travelled by solute} / \text{Distance travelled by solvent}$$

### 2.6 Total Phenolic Content (TPC)

The Folin-Ciocalteu method [33] was used for the determination of TPC. 100 µl of different concentrations extracts was mixed with 500 µl of water and 100 µl of Folin-Ciocalteu reagent. The mixture was incubated for six minutes. Further, 500µl of distilled water and 1 ml of 7% sodium carbonate was added to the reaction mixture. The absorbance was taken after ninety minutes at 760 nm spectrophotometrically. The gallic acid equivalents (mg GAE/g) were used to calculate the TPC. Each test was carried out in triplicates.

## 2.7 Total Flavonoid Content (TFC)

The aluminum chloride complex formation assay, which was performed by Piyanete et al. [34], was used to determine the TFC of the methanolic extract of *A. marmelos* leaves. "The quercetin was utilized as a reference, the flavonoid content was calculated as mg QE/g. After mixing 100  $\mu$ l of the quercetin dilution with 500  $\mu$ l of distilled water and 100  $\mu$ l of 5% sodium nitrate, the mixture was left to stand for 6 minutes. Subsequently, 200  $\mu$ l of 1M NaOH solution was added after 150  $\mu$ l of a 10%  $AlCl_3$  solution had been added and incubated for 5 minutes. This reaction mixture's absorbance was measured at 510 nm. All the procedures were performed in triplicates" [32].

## 2.8 Nitric Oxide Radical Scavenging Assay

"Free radicals generated from sodium nitroprusside (SNP) were measured according to the earlier described method [31] with some modifications. Different concentration of reaction mixture containing SNP (15 mM) in PBS (pH 7.3) with and without sample, incubated at 25°C for 210 mins. After adding the Griess reagent, the reaction mixture was incubated for 10 minutes at room temperature. The standard used was ascorbic acid. The absorbance was measured at 560 nm using a UV-Vis microplate reader" [32].

## 2.9 Superoxide Anion Scavenging Assay

"Based on a previously described method [35] with minor modifications, the reduction of NBT determined the total antioxidant capacity of methanolic extract. Phosphate buffer (20 mM, pH 7.4), PMS (60 $\mu$ M), NBT (156 $\mu$ M), and different conc. of extract were all included in the 1-mL reaction mixture. After incubation for 5 min at 25°C temperature, the absorbance was taken at 560 nm against an appropriate blank solution. BHT was used as reference control" [32].

## 2.10 (GC-MS) Analysis

The GC-MS analysis of the methanolic extract of *A. marmelos* leaves was carried out on a Perkin Elmer Turbo Mass Spectrophotometer using a Perkin Elmer autosampler XLGC. The column was a Perkin Elmer Elite - 5 capillary columns with a film thickness of 0.25 mm and a length of 30 m. It was made of 95% dimethylpolysiloxane. At a flow rate of 0.5 ml/min, helium (99.999 percent) was used as the carrier gas. An

injection length of one liter was employed. The GC's inlet temperature was kept at 250°C, and it had a programmed oven temperature of 110°C (isothermal for two minutes). After that, it increased by 10°C/min to 200°C, then by 5°C/min to 280°C, and finally by 5°C/min to 280°C, with a 5°C/min isothermal at 280°C. The GC ran in 30 minutes. The source temperature is maintained at 180°C, and the temperature of the MS transfer line is maintained at 200°C. Total Ion Count was utilized for data 60 Sharma and Kumar evaluation of compound detection and quantification (TIC), and electron impact ionization at 70eV was used for the GC-MS analysis. The spectrum of the components was compared to the known components stored in the GC-MS library. For peak area measurement and data processing, Turbo-Mass OCPTVS-Demo SPL programme 19 was used.

## 2.11 Statistical Analysis

The results were shown as mean  $\pm$  SEM. The student's t test (paired) was used at the 5% level to determine the significance of the observed differences. The comparisons were done between reference controls and methanolic leaves extracts of *A. marmelos*.

## 3. RESULTS


Both the humans and animals require secondary metabolites. The biochemical tests showed (Table 1) the presence of different secondary metabolites in the aqueous and methanolic extracts. The results depend on the solvent as well as the qualitative detection methods used. It was observed that there are not many differences in the presence or absence of secondary metabolites when 5 & 10 gm of plant material were used for aqueous extraction. For the determination of secondary metabolites, aqueous and methanolic extract are equally important. The total 5 spots with different Rf values (0.31, 0.59, 0.63, 0.71 and 0.79) were detected by TLC in the methanolic extract of *A. marmelos* leaves (Fig. 1). Here we used alanine amino acid as a standard and its Rf value was 0.69.

Ascorbic acid showed 91.0% scavenging activity whereas methanolic extract was found to be 64.40% at a concentration of 1000 mg/ml (Fig. 2). Significant differences have been observed in the antioxidant potential between the ascorbic acid (reference control) and methanolic extract of *A. marmelos* ( $p= 0.00985$ ).

**Table 1. Qualitative phytochemical analysis of the aqueous and methanolic extracts of *A. marmelos* leaves [32]**

SNo.	Phytochemical Tests	Aqueous Extract		Methanolic Extract
		10 gm	5 gm	
1.	<b>Alkaloids</b>			
	Mayer's	+ ve	+ ve	+ ve
	Wagner's	+ ve	- ve	+ve
	Hager's	+ ve	+ ve	+ ve
2.	<b>Carbohydrates</b>			
	Molisch	- ve	+ ve	- ve
	Barfoed's	- ve	+ ve	- ve
3.	<b>Reducing Sugars</b>			
	Fehling's	+ ve	+ ve	+ ve
	Benedict's	+ ve	+ ve	+ ve
4.	<b>Flavonoids</b>			
	Alkaline Reagent	+ ve	-ve	-ve
	Lead Acetate	+ ve	+ ve	+ ve
5.	<b>Glycosides</b>			
	Borntrager's	+ ve	+ ve	+ ve
	Legal's	- ve	- ve	-ve
	Keller-killiani	- ve	- ve	+ ve
6.	<b>Tannin &amp; phenolic</b>			
	Ferric Chloride	+ ve	+ ve	+ ve
	Lead Acetate	+ ve	+ ve	+ ve
	Dilute iodine solution	+ ve	+ ve	+ ve
7.	<b>Saponin</b>			
	Froth	-ve	-ve	+ve
8.	<b>Protein &amp; A.A.</b>			
	Ninhydrin	+ ve	+ ve	+ve
	Biuret	-ve	-ve	-ve
9.	<b>Triterpenoids &amp; Steroids</b>			
	Salkowski's	-ve	-ve	+ ve
10.	<b>Hydrolysable tannin</b>	+ ve	+ ve	+ ve

(+) indicates presence while, (-) indicates the absence of the components



**Fig. 1. TLC plate showing spots having different Rf values (0.79, 0.71, 0.63, 0.59 & 0.31) of methanolic extract of *A. marmelos* leaves**




Fig. 2. The Antioxidant activity of methanolic extract of *A. marmelous* leaves

Table 2. Total Flavonoid & phenolic content of methanolic extract of *A. marmelous* leaves

Total Flavonoid Content (mgQE/g)		Total Phenolic Content (mgGAE/g)	
Conc. (µg/ml)	<i>A. marmelous</i>	Conc. (µg/ml)	<i>A. marmelous</i>
1000	123±13.5	150	50±5.3
500	128±15.6	120	40±4.1
250	112±11.7	90	27±2.4
125	392±21.6	60	15±2.2
62.5	784±26.8	30	
<b>Mean ± SD</b>	<b>307.8 ± 130.12</b>	<b>Mean ± SD</b>	<b>33± 7.62</b>




Fig. 3. Nitric oxide scavenging activity of methanolic extract of *A. marmelous* leaves

The antioxidant potential observed was due to the presence of phenolic and flavonoid contents. The mean values of TPC and TFC were  $33 \pm 7.62$  mg GAE/g and  $307.8 \pm 130.12$  mg QE/g respectively (Table 2).

The dose dependent NO and superoxide radical scavenging activity of the methanolic extract of *A. marmelous* leaves were observed (Figs. 3 and 4). Significant differences have been observed in the NO and superoxide radical scavenging

activity between the ascorbic acid (reference control) & methanolic extract of *A. marmelos*, respectively ( $p= 0.000161$ ;  $p=0.036419$ ). The active principle analyzed by GC-MS with MW, MF, peak area%, RT, and composition of the bioactive components of methanolic extract of *A. marmelos* leaves (Table 3). The GC-MS chromatogram of the identified compounds is shown in (Fig. 5).

Many components that are biologically active were observed. The area with highest peak of Perylo[1,12-def]-1,3-dioxepin-5,11-dione,6,12-dihydroxy-8,9-bis(2-hydroxypropyl)-7,10-dimethoxy-, tereoisomer (3.502%), Lutein (3.040%), Rhodopin (3.040%), Hyocholic acid (2.468%) and Hematoporphyrin (1.311%) shown different biological activities like Antioxidant, anticancerous, antileishmanial and antimalarial activities etc.

#### 4. DISCUSSION

*A. marmelos* have been traditionally used for many ailments and the nutritional benefits are also reported. The best solvents for extracting this plant's metabolites are methanol and water, with ethanol coming in second [41]. The phytochemistry of *A. marmelos* has been thoroughly investigated, and it has been

discovered that the plant contains a wide range of physiologically active substances. *A. marmelos* contains a number of important phytochemicals, such as: Alkaloids which are well-known for their medicinal effects. Marmelosin, aegeline, and marmesin are among the alkaloids that have been found in leaves & roots of *A. marmelos* [42]. *A. marmelos* fruit contain abundance of tannin, and studies have indicated that tannins have potent anti-inflammatory and antioxidant properties. Our results also showed the presence of tannins in the aqueous and methanolic leaf extracts. Flavonoids are a class of chemicals that have antioxidant, anti-cancer, and anti-inflammatory properties. Roots and leaves of *A. marmelos* contain flavonoids, which are recognized for their anti-cancer, antioxidant, and anti-inflammatory properties [43]. *A. marmelos* has been shown to contain terpenoids, some of which have been demonstrated to have antibacterial & antifungal properties. Saponins are also present in the leaves and fruit of *A. marmelos*, which are recognized for their ability to foam and emulsify. Studies have demonstrated the anti-inflammatory & antinociceptive effects of several of these chemicals [44]. Glycosides are well-known for their medicinal properties and have been shown to have antinociceptive and anti-inflammatory activities [45].




Fig. 4. The superoxide radical scavenging activity of methanolic extract of *A. marmelos* leaves




Fig. 5. Total Ion Chromatogram of methanolic extract of *A.marmelous* leaves



**Table 3. Compounds identified by GC-MS analysis of methanolic extract of *A. marmelous* leaves**

<b>Sr. No</b>	<b>RT</b>	<b>PP%</b>	<b>Compound name</b>	<b>M.F.</b>	<b>M.W.</b>	<b>Biological Activities**</b>
1.	20.610	1.311	Hematoporphyrin	C <sub>34</sub> H <sub>38</sub> N <sub>4</sub> O <sub>6</sub>	598.7g/mol	Phototherapy of malignant neoplasms [36]
2.	20.751	2.468	Hyocholic acid	C <sub>24</sub> H <sub>40</sub> O <sub>5</sub>	408.6g/mol	Mouse metabolite, a human urinary metabolite and a rat metabolite Antiplasmodial activity antitrypanosomal activity [37]
3.	23.178	3.502	Perylo[1,12-def]-1,3-dioxepin-5,11-dione, 6,12-dihydroxy-8,9-bis(2-hydroxypropyl)-7,10-dimethoxy-, stereoisomer	C <sub>29</sub> H <sub>26</sub> O <sub>10</sub>	534.5g/mol	Antimalarial Antileishmanial [38]
4.	28.321	3.040	Lutein	C <sub>40</sub> H <sub>56</sub> O <sub>2</sub>	568.9g/mol	An antioxidant Plant metabolite Antiviral activity [39]
5.	28.321	3.040	Rhodopin	C <sub>40</sub> H <sub>58</sub> O	554.9g/mol	Bacterial metabolite [40]

\*\* Biological Activities Based on the PubChem Database

In synthetic chemistry, TLC is widely used method for component identification, purity analysis, and progress of reaction. Additionally, it enables the solvent system to be optimized for a specific separation problem. It is considerably faster than column chromatography and just needs a minimal amount of a compound. A thin, homogeneous coating of silica gel or alumina is applied to a piece of glass, metal, or rigid plastic in TLC. A suitable liquid solvent or solvent mixture serves as the mobile phase [46].

Antioxidants are very important in preventing human illnesses by acting as scavengers of free radicals, pro-oxidant metal complexes, reducing agents, and quenchers of singlet oxygen production [47]. Because of their ability to scavenge, antioxidants are able to manage certain conditions, including as AIDS, cancer, heart disease, neurological disorders, and cataracts [48]. Scientific knowledge regarding the antioxidant abilities of many plants is currently insufficient, particularly those less commonly used in food and medicine. Thus, evaluating these characteristics continues to be a worthwhile and fascinating endeavor, especially when it comes to discovering fresh sources of functional foods, natural antioxidants, and nutraceuticals. It has been suggested that synthetic antioxidants such as gallic acid esters, BHT (butylated hydroxyl toluene), tertiary butylated hydroxy quinone, and BHA (butylated hydroxy anisole) are carcinogenic. As a result, there are strict limitations on their use, and replacing them with antioxidants that occur naturally is recommended. These synthetic antioxidants likewise exhibit minimal antioxidant activity and poor solubility [49]. Finding a more potent antioxidant with less adverse effects from natural sources is still necessary. It has been shown that plants containing flavonoids exhibit antioxidant effects [50].

Using the phosphomolybdenum assay, bael leaf's antioxidant activity was evaluated. It is widely known that plant flavonoids and phenolics in general are potent antioxidants and scavengers of free radicals [51,52]. The majority of plants' capacity to scavenge radicals has been attributed to substances like flavonoids, which include hydroxyls [53]. Leaves of *A. marmelos* were found to contain, phenolic compounds, flavonoids, fats, triterpenoids, glycosides, carbohydrate and alkaloids as their chemical constituents. Thus, the current investigation has

confirmed that the leaves of remedial plants may be a rich source of antioxidants. The substantial amount of phenolic hydroxyl groups that phenolics possess may account for their potent ability to scavenge free radicals.

## 5. CONCLUSION

The findings indicate that *Aegle marmelos* leaves have many phytoconstituents and strong antioxidant properties and that their presence or absence depends on the solvent and the methodology used. The present study reveals that methanolic extract contains more phytochemicals compare to aqueous extract, it is enough excited to explore more information about this medicinal plant, therefore it inhibits production of free radicals. GC-MS analysis revealed many compounds viz. Hematoporphyrin, Hyocholic acid, Lutein Rhodopin, etc. with known biological activities. In addition, the current study suggests the possibility of manufacturing health supplements using these medicinal plants. In many respects, *A. marmelos* is a unique source of drugs. Each and every part of this plant treat like a medicine and used by human beings and for their cattles also.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

The author(s) thus declare that during the drafting or editing of manuscripts, NO generative AI tools, such as large language models (ChatGPT, COPILOT, etc.), or text-to-image generators, were employed.

## DATA AVAILABILITY STATEMENT

Data set supporting this manuscript is included within the manuscript. Any further information should be requested from the corresponding author.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

It is not applicable.

## ACKNOWLEDGEMENTS

The authors are grateful to the Innovation Centre, Bundelkhand University, Jhansi for the GC-MS analysis.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Makkar HPS, Norvsambuu T, Lkhavatsere S and Becker K. *Tropicultura*.2009;27:159-167.
- Akindele AJ, Adeyemi. Anti-inflammatory activity of the aqueous leaf extracts of *Byrsocarpus coccineus*. *J. Fitoterapia*. 2007;78:25-28.
- Salah N, Miller NJ, Paganga G, Tigburg L, Bolwell GP and Rice- Evans CA. *Archives of Biochemistry and Biophysics*. 1995;322:339-346.
- Edge R, McGarvey DJ and Truscott TG. The Carotenoids as Anti-Oxidants: A Review. *Journal of Photochemistry and Photobiology B*. 1997;41(3):189-200.
- Papas AM. In phytochemicals in Nutrition and Health, Meskin, M. S., Bidlack, W. R., Davies, A. J. and Omaye. S. T. New York: CRC Press, 2002;61-78.
- Aruoma OL, Halliwell B, Aeschbach R and Loligers J. Antioxidant and pro-oxidant properties of active rosemary constituents: Carnosol and Carnosic acid. *Xenobiotica*. 1992;22:257-268.
- Poljsak B, Šuput D, Milisav I. Achieving the balance between ROS and antioxidants: when to use the synthetic antioxidants. *Oxid. Med. Cell. Longev*; 2013.
- Antolovich M, Prenzler PD, Patsalides E, McDonald S, Robards K. Methods for testing antioxidant activity. *Analyst*, 2002;127(1):183-198.
- Shenoy R, Shirwaikar A. Anti-inflammatory and free radical scavenging studies of *Hyptissuaveolens* (Labiatae). *Indian Drugs*. 2002;39(11):574-577.
- Alam MN, Bristi NJ, Rafiquzzaman M. Review on *in vivo* and *in vitro* methods evaluation of antioxidant activity. *Saudi Pharmaceutical Journal*. 2013;21(2):143-152.
- Kinnula VL, Crapo JD. Superoxide dismutases in malignant cells and human tumors. *Free Radic. Biol. Med*. 2004;36(6):718-744.
- Singh U, Jialal I. Oxidative stress and atherosclerosis. *Pathophysiology*. 2006;13(3):129-142.
- Sas K, Robotka H, Toldi J, Vécsei L. Mitochondria metabolic disturbances, oxidative stress and the kynurenine system, with focus on neurodegenerative disorders. *J. Neurol. Sci*. 2007;257(1):221-239.
- Smith MA, Rottkamp CA, Nunomura A, Raina AK, Perry G. Oxidative stress in Alzheimer's disease. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*, 2000;1502(1):139-144.
- Guidi I, Galimberti D, Lonati S, Novembrino C, Bamonti F, Tiriticco M, Fenoglio C, Venturelli E, Baron P, Bresolin N, Scarpini E. Oxidative imbalance in patients with mild cognitive impairment and Alzheimer's disease. *Neurobiol. Aging*. 2006;27(2):262-269.
- Bolton JL, Trush MA, Penning TM, Dryhurst G, Monks TJ. Role of quinones in toxicology. *Chem. Res. Toxicol*. 2000;13(3):135-160.
- Arteel GE. Oxidants and antioxidants in alcohol-induced liver disease. *Gastroenterology*. 2003;124(3):778-790.
- Ramakrishna B, Varghese R, Jayakumar S, Mathan M, Balasubramanian K. Circulating antioxidants in ulcerative colitis and their relationship to disease severity and activity. *J. Gastroenterol. Hepatol*. 1997;12(7):490-494.
- Upston JM, Kritharides L, Stocker R. The role of vitamin E in atherosclerosis. *Prog. Lipid Res*. 2003;42(5):405-422.
- Hyun D-H, Hernandez JO, Mattson MP, de Cabo R. The plasma membrane redox system in aging. *Ageing Research Reviews*. 2006;5(2):209-220.
- Lee KW and Lee HJ. The roles of polyphenols in cancer chemoprevention. *Biofactors*. 2006;26(2):105-121.
- Moon Y, Wang X and Morris M. Dietary flavonoids: Effects on xenobiotic and carcinogen metabolism. *Toxicol. In Vitro*. 2006;20(2):187-210
- Valko M, Izakovic M, Mazur M, Rhodes CJ and Telser J. Role of oxygen radicals in DNA damage and cancer incidence. *Mol. Cell. Biochem*. 2004;266(1-2):37-56.
- Poulsen HE, Prieme H and Loft S. Role of oxidative DNA damage in cancer initiation and promotion. *Eur. J. Cancer Prev*. 1998;7(1):9-16.
- Laranjinha JA, Almeida LM and Madeira VM. Reactivity of dietary phenolic acids with peroxy radicals: Antioxidant activity upon low density lipoprotein peroxidation. *Biochem. Pharmacol*. 1994;48(3):487-94.

26. Gropper SS, Smith JL and Groff JL. Advanced nutrition and human metabolism. Cengage Learning, Wadsworth; 2009.
27. Cook NC, Samman S. Flavonoids-Chemistry, metabolism, cardioprotective effects, and dietary sources. Nutrition. Biochemistry. 1996;7:66-76.
28. Singh V, Kumar R. Study of Phytochemical Analysis and Antioxidant Activity of *Allium sativum* of Bundelkhand Region. Int. J. Life. Sci. Scienti. Res. 2017;3(6):1451-1458.
29. Bisht D, Prakash D, Kumar R, Shakya AK, Shrivastava S. Phytochemical profiling and nephroprotective potential of ethanolic leaf extract of *Polyalthia longifolia* against cisplatin-induced oxidative stress in rat model. Journal of Ethnopharmacology. 2024;326:117922
30. Sharma S, Yadav S, Kumar R. GC-MS Analysis, Antioxidant and Free Radical Scavenging Activity of *Embllica officinalis* Gaertn (Amla) Fruit Extracts. Bull. Env.Pharmacol. Life Sci. 2024;13(3):18-26
31. Prieto P, Pineda M, Anguilar M. Spectrophotometric quantitation of antioxidant capacity through the formation of a Phosphomolybdenum Complex: Specific application to the determination of Vitamin E. Anal. Biochem. 1999;269:337-341.
32. Sharma S, Kumar R. Free radical scavenging activity and gc-ms analysis of methanolic extract of *Ficus racemosa* L. Leaves of bundelkhand region. Journal of Phytological Research. 2021 Jan 1;34(1).
33. Odabasoglu F, Aslan A, Cakir A, Suleyman H, Karagoz Y, Halici M, et al. Comparison of antioxidant activity and phenolic content of three lichen species. Phytotherapy Research. 2004;18(11):938-941.
34. Piyanete C, Meechai P, Nakbanpottecc W. Antioxidant activities and phenolic contents of extracts from *Salvinia olesta* and *Eichornia crassipes*. Res J Biol Sci., 2009;4:1113-1117.
35. Marcocci L, Packer L, Droy-Lefaix MT, Sekaki A and Gardes-Albert M. Antioxidant action of Ginkgobiloba extracts EGB 761. Methods Enzymol. 1994;234:462-475.
36. National Center for Biotechnology Information. PubChem Compound Summary for CID 11103, Hematoporphyrin; 2021. Retrieved January 25, 2021.
37. National Center for Biotechnology Information. PubChem Compound Summary for CID 92805, Hyocholic acid. Retrieved January 25, 2021. Available: <https://pubchem.ncbi.nlm.nih.gov/compound/Hematoporphyrin>.
38. National Center for Biotechnology Information. PubChem Compound Summary for CID 91617, Cercosporin. Retrieved January 25, 2021. Available: <https://pubchem.ncbi.nlm.nih.gov/compound/Hyocholic-acid>.
39. National Center for Biotechnology Information. PubChem Compound Summary for CID 5281243, Lutein. Retrieved January 25, 2021. Available: <https://pubchem.ncbi.nlm.nih.gov/compound/Cercosporin>.
40. National Center for Biotechnology Information. PubChem Compound Summary for CID 5365880, Rhodopin. Retrieved January 25, 2021. Available: <https://pubchem.ncbi.nlm.nih.gov/compound/Lutein>.
41. Veer B, Singh R. Phytochemical Screening and Antioxidant Activities of *Aegle marmelos* Leaves. Anal. Chem. Lett. 2019;9(4):478-485.
42. Baliga MS, Bhat HP, Joseph N, Fazal F. Phytochemistry and medicinal uses of the bael fruit (*Aegle marmelos* Correa): a concise review. Food Res. Int. 2011;44(7):1768-1775.
43. Choudhary Y, Saxena A, Kumar Y, Kumar S, Pratap V. Phytochemistry, pharmacological and traditional uses of *Aegle marmelos*. Pharm. Biosci. Journal. 2017;20:27-33.
44. Sharma B, Sharma P. Constituents of *Aegle marmelos*. Planta Med. 1981;43(09):102-103.
45. Dhankhar S, Ruhil S, Balhara M, Dhankhar S, Chhillar AK. *Aegle marmelos* (Linn.) Correa: A potential source of Phytomedicine. J. Med.Plants Res. 2011;5(9):1497-1507.
46. Stoddard JM, Nguyen L, Mata-Chavez H and Nguyen K. TLC plates as a convenient platform for solvent-free reactions Chemical Communications (Camb). 2007;28(12):1240-1241.
47. Andlauer W, Furst P: Antioxidative power of phytochemicals with special reference to cereales. Cereal Foods World. 1998; 43:356-359.

48. Middleton E, Kandaswami C, Theoharides TC: The effects of plants flavonoid on mammalian cell: Implication for inflammation, heart disease, and cancer. *Pharmacological Review*.2000;52: 673-751.
49. Barlow SM. Toxicological aspects of antioxidants used as food additives. *In Food Antioxidants*, Hudson BJB (ed.) Elsevier, London. 1990;253-307.
50. Cook NC, Samman S: Flavonoids-chemistry, metabolism, cardioprotective effects, and dietary sources. *Nutritional Biochemistry*. 1996;7:66-76.
51. Shahidi F, Wanasundara JPD. Phenolic antioxidants. *Critical Reviews in Food Science and Nutrition*. 1992;32:67–103.
52. Sofowora A. Medicinal plants and traditional medicine in Africa, Spectrum books, Ind, Ibadan, Nigeria. 1993;289.
53. Das NP, Pereira TA. Effects of flavonoids on thermal autoxidation of palm oil: Structure-activity relationship. *J Am Oil Chem Soc*. 1990;67:255-258.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/124581>