



# **The Effect of Aromatherapy with *Citrus aurantium* on Anxiety during MRI Imaging in Patients with Spinal Disorders: A Randomized Clinical Trial**

**Nishtman Ahani<sup>1</sup>, Kamal Salehi<sup>2\*</sup>, Jamal Seidi<sup>2</sup>, Bakhtyar Salehi<sup>3</sup> and Bijan Nouri<sup>4</sup>**

<sup>1</sup>Student Research Committee, Kurdistan University of Medical Science, Sanandaj, Iran.

<sup>2</sup>Clinical Care Research Center, Kurdistan University of Medical Sciences, Sanandaj, Iran.

<sup>3</sup>MRI Department, Imam Khomeini Hospital of Saqqez, Iran.

<sup>4</sup>Social Determinants of Health Research Center, Research Institute for Health Development, Kurdistan University of Medical Sciences, Sanandaj, Iran.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/JPRI/2019/v26i230134

### Editor(s):

(1) Dr. Jinyong Peng, Professor, College of Pharmacy, Dalian Medical University, Dalian, China.

### Reviewers:

(1) Vijaya Krishnan, (PT) MGM College of Physiotherapy, Navi Mumbai, India.

(2) Daniela Benedec, "Iuliu Hatieganu" University of Medicine and Pharmacy Cluj-Napoca, Romania.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/46877>

**Original Research Article**

**Received 24 October 2018**

**Accepted 26 January 2019**

**Published 27 March 2019**

## **ABSTRACT**

**Objective:** Anxiety is one of the most significant problems of patients during magnetic resonance imaging (MRI). Anxiety affects the quality of images, increased costs and repeating the imaging process. Aromatherapy is regarded a complementary therapy that is assumed to be highly effective in reducing anxiety. The present study aims at investigating the effect of aromatherapy with *Citrus aurantium* on anxiety in MRI imaging.

**Materials and Methods:** This randomized clinical trial was done on 80 patients with spinal disorders that were randomly assigned to the intervention and control groups. In the intervention group, inhaling *Citrus aurantium* essential oil was conducted two minutes before entering the MRI machine and right before entering the MRI machine. As for the control group, the patients were provided with only the common cares of the medical center. Right after the imaging process, the

\*Corresponding author: E-mail: salehikamal88@gmail.com;

anxiety experienced was evaluated based on the Magnetic Resonance Imaging-Anxiety Questionnaire (MRI-AQ). The data collected were analyzed in SPSS-21 with statistical tests including independent T-test, Fisher's exact test, and chi-squared test.

**Results:** The samples of both groups were homogenous in terms of age, sex, marital status, educational level, occupation, and the kind of disease. The mean of MRI anxiety experience was  $23.32 \pm 12.72$  and  $37.4 \pm 15.41$  in the intervention and control group respectively; the independent T-test indicated a significant difference.

**Conclusion:** *Citrus aurantium* is likely to reduce the anxiety in MRI imaging. Given the proper effectiveness, inexpensiveness, and the easy administration of this aromatherapy, it is recommended to be applied for reducing the MRI anxiety.

**Keywords:** Aromatherapy; *Citrus aurantium*; randomized clinical trial; anxiety; MRI.

## 1. INTRODUCTION

Magnetic Resonance Imaging (MRI) is a comprehensive imaging method [1] and a main diagnostic tool in patients with spinal cord disorders and abnormalities [2]. Although MRI is physically safe, anxiety is the most important problem the patients deal with [1]. In their study, Thu et al. [3] indicated that 25% of patients suffer from moderate to severe anxiety during the scanning process [3]. Chapman maintains that the scanning experience is likely to create an anxiety level similar to that of the operation [4]. Imaging anxiety can increase the patient's movement during the imaging; this results in motor noise, reduced quality of the images, and reduced diagnostic value [5]. In severe cases, the imaging is likely to be interrupted and the patient avoids continuing it [6]. Annually, as many as 2 million scans are interrupted owing to anxiety and the interruption of imaging; this imposes considerable costs on the medical system [7].

Various methods are conducted for reducing MRI anxiety. In some cases, Benzodiazepines are applied for controlling anxiety; they are likely to result in the patient's drowsiness and his/her fall. Thus, there is an increased tendency towards applying non-medicinal methods for reducing anxiety [8]. Various methods have been conducted about non-medicinal methods. Tazegul et al. [9] reduced the scanning anxiety through providing information and communications with the patients via intercom [9]. In their study, Chapman et al. [4] indicated that an MRI simulator (before the scanning) can reduce the confusion and anxiety of MRI machine. In another study, Mohammad Kamal et al (2013) reduced the patients' anxiety through providing information, creating enjoyable posters, presenting pamphlets in the waiting room, training stress coping strategies and proper communicative methods with nurses of the

scanning department, and counselling on cognitive strategies [10]. The non-medicinal methods include aromatherapy, massage therapy, muscle relaxation, music therapy, and guided visualization [11].

Aromatherapy is a complementary therapy that is assumed to be effective in reducing anxiety, fatigue, insomnia, depression [12], and pain [13]. In aromatherapy, the essential oil is likely to increase the individual's increased feeling of calm and peace by affecting the limbic system of the brain [14]. *Citrus aurantium* essential oil is one the most commonly used aromas in aromatherapy [15]. *Citrus aurantium* is one of the highly used medicinal plants. *Citrus aurantium* is a local plant that grows in the north and south of Iran. This plant is applied as a sedative, hypnotic, appetizer, and heart palpitation remedy. The components of *Citrus aurantium* extract include alkaloids, linalool, linalyl acetate, limonene, lymonoyds, flavonoids, msyryn [12]. Moreover, flavonoids function as the agonist of benzodiazepines, and they reduce the patient's anxiety in this way [16]. In their study, Leite et al. [17] indicated that applying *Citrus aurantium* through inhaling has anti-anxiety effects [17]. In another study, Namazi et al. [16] indicated that the essential oil of *Citrus aurantium* results in reduced anxiety during the first stage of labor [16].

Since MRI is a diagnostic method with special features including high diagnostic value and high prevalence and since it is time consuming, anxiety is one of its most significant problems. Reducing the patients' anxiety in this diagnostic method seems to be highly effective. At present, numerous clinical studies are being conducted around the world on different applications of aromatherapy and using herbal essential oils. It seems that the essential oil of *Citrus aurantium* can be effective as an inexpensive, simple, and safe method for reducing anxiety in MRI. Thus,

the present study was conducted to determine the effect of aromatherapy with *Citrus aurantium* on the anxiety experienced during MRI.

## 2. MATERIALS AND METHODS

The present study is an After-Only with Control Group clinical trial conducted in 2018 on 80 patients with spinal disorders. They had referred to the MRI department of Imam Khomeini Hospital of Saqqez affiliated with Kurdistan University of Medical Sciences. The study was conducted after being approved by the Research Council of Kurdistan University of Medical Sciences, acquiring confirmation from the Ethics Committee, and being registered at the Iranian Registry of Clinical Trials with the code of IRCT201801170338414N1.

The inclusion criteria of the present study are as follows: age of 18-70 years; not suffering from a specific disease (allergy, olfactory disorder, and anxiety disorder); no addiction to psychedelic drugs; lack of respiratory problems; lack of experiencing MRI scan in the past; outpatient patients; spinal disorders (neck and waist); spinal abnormalities (Scoliosis, Kyphosis, Lordosis); having minor or moderate anxiety according to Spielberger's Test Anxiety Inventory in both groups of control and intervention. The exclusion criteria are: any medical or surgical problem requiring immediate medical intervention during the study; sensitivity to the essential oil of *Citrus aurantium*; dissatisfaction after inhaling the essential oil; the sample's voluntary renouncement to continue his/her participation for any reason; having severe anxiety; suffering from generalized anxiety disorder; suffering from Obsessive Compulsive Disorder; having claustrophobia; and suffering from Social Phobia Disorder.

The statistical population of the present study include all patients registered at the MRI department. With the interval confidence of 95% and the test power of 80%, the number of samples was estimated to be 40 for each group through applying the sample size formula. For sampling, the simple random method was applied. After numbering the patients' names, a table of random number was used for the initial sampling. After conducting the sampling, the samples were re-divided into two groups of control and *Citrus aurantium* by applying a simple random method. It is also worth noting that, the experiment was conducted weekly and separately for different groups, i.e. in one week

the MRI of *Citrus aurantium* was conducted and in the following week, the MRI of the control group was administered; this measure was taken so that the control group patients were not unwantedly affected by the essential oil of *Citrus aurantium*. Before initiating the intervention, the patients' anxiety was evaluated according to Spielberger's Test Anxiety Inventory. The patients with minor and moderate anxiety were entered into the study, and the patients with severe anxiety were excluded from the study. The essential oil *Citrus aurantium* required for conducting the study was bought from a reputable company in Tehran. After determining the concentration with a graviometry method (conducted by the School of Pharmacy of Tehran University of Medical Sciences), the essential oil was used. This essence had 10 mg of *Citrus aurantium* essence provided in 100 ml.

The intervention of the present study was conducted by inhaling the essential oil of *Citrus aurantium* on a 2×2 gauze impregnated with 3 drops of *Citrus aurantium* 10% attached to the patient's collar at two intervals i.e. 10 minutes before entering the MRI machine and right before entering the MRI machine. Right after the imaging process, the anxiety experienced was evaluated based on the Magnetic Resonance Imaging-Anxiety Questionnaire (MRI-AQ). As for the control group, the patients were provided only with the common cares of the medical center. The procedure was conducted until all research samples underwent the process. The patients' heartbeat, blood pressure, nausea, vomiting, and headache were controlled after administering the aromatherapy. None of the patients relinquished during the process of the study.

The data collection tool of the present study is divided into three parts. The first part includes the individual characteristics (age, sex, educational level, marital status, economic status, and the type of the scan conducted). The second part was Spielberger's anxiety questionnaire that included 20 questions; it was divided into three anxiety groups including minor (scores of 20-29), moderate (scores of 40-59), and severe (60-80) [18]. This questionnaire is widely used in clinical studies to measure anxiety; the correlation coefficient acquired for this test is 0.85-0.91 [19]. In Iran, the reliability coefficient of this questionnaire has been separately investigated in two groups of norm and criterion. The reliability of norm group (600) on an overt and covert anxiety scale (based on

Cronbach's alpha) was 0.9084 and 0.9025 respectively. This value was 0.9418 in the criterion group [20]. The third part was MRI-AQ including 15 items. Each item had 4 options including 1. Never, 2. To some extent, 3. To an average level, and 4. Too much. This questionnaire describes the patient's mental states and anxiety during the imaging. This questionnaire was standardized in 2015 by Ahlander et al. [21] for investigating the anxiety of MRI [21]. For determining the validity of the tool, it was first translated into Persian and the items were adapted to the clinical and social conditions existing in Iran. For modification and correction, the tool was then presented to 10 faculty members of Nursing and Midwifery Faculty of Kurdistan University of Medical Sciences. After collecting information and conducting the required corrections and modifications, the questionnaire was presented to the Ethics Committee of the university. At the end, the modified version was applied. For determining the reliability of the questionnaire, the validated version was presented to 40 patients who had voluntarily referred to the MRI department of Saqqez Hospital. The items were then evaluated by using Cronbach's alpha. The reliability score was 0.84. For analyzing the data of the present study, SPSS-21 was applied. First, mean, standard deviation, and the interval confidence of 95% were evaluated in both groups. For investigating the homogeneity of both groups in terms of background variables, Fisher's exact test and chi-squared test were used. Then, for investigating the effect of the intervention, parametric Independent-T test was applied.

The intended ethical considerations of the present study are explaining the purpose and nature of the study for all samples, acquiring informed consent from the samples, assuring the patients about the confidentiality of the data, telling the patients that there is no need to write their first and last names in the questionnaire, and providing the samples and officials with the summary of the research findings.

### 3. RESULTS

The patients investigated were in the age range of 18-65 years with the mean age of 30.36 and standard deviation of 15.73. Most of the participants of the present study were less than 30 years old (35%), female (55%), married (65%), self-employed (40%), and suffered from

discopathy (95%). The data analysis indicated that the intervention and control group were homogenous in terms of demographic features including age, sex, marital status, educational level, occupation, and kind of spinal disorder; they were not significantly different ( $p < 0.05$ ) (Table 1).

The mean anxiety experienced during MRI was  $23.32 \pm 12.72$  in the intervention group and  $37.4 \pm 15.41$  in the control group. The independent T-test indicated a significant difference between the mean of anxiety in intervention and control groups ( $P < 0.05$ ).

### 4. DISCUSSION

The findings of the present study indicated that aromatherapy with *Citrus aurantium* is likely to reduce MRI anxiety.

The findings of the present study are consistent with those of the studies conducted by Sharifi Pour et al. [22] and Akhlaghi et al. [23] in terms of the aromatherapy effectiveness on reducing anxiety. Sharifi Pour et al, indicated that the essence of *Citrus aurantium* is effective in reducing the post-operational anxiety in caesarean patients [22]. In the study conducted by Sharifi Pour et al, the severity of anxiety was measured by using Spielberger's anxiety tool in two stages: right after the operation and 12 hours after operation. The control group underwent a test with placebo. However, in the present study, the severity of anxiety was measured only at one stage (after the intervention) with the Resonance Imaging-Anxiety Questionnaire (MRI-AQ). Moreover, in their study, Akhlaghi et al. [23], indicated that *Citrus aurantium* reduces pre-operation anxiety [23]; this is consistent with the findings of the present study. For reducing MRI anxiety, some other methods have been applied as well. In their study, Chapman et al. [4], indicated that an MRI simulator (before the scanning) can reduce the confusion and anxiety of MRI machine. In another study, Mohammad Kamal et al, reduced the patients' anxiety through providing information, creating enjoyable posters, presenting pamphlets in the waiting room, training stress coping strategies and proper communicative methods with nurses of the scanning department, and counselling on cognitive strategies [10]. Tazegul et al. [9], reduced the scanning anxiety through providing information and communications with the patients via intercom [9]. Tugwell et al. [24]

**Table 1. Table of initial indices in the intervention and control group**

Variable	Control		Intervention		P value
	Number	Percentage	Number	Percentage	
<b>Age group</b>					
20- 30	14	35	18	45	Fisher's test 0.44
31-40	13	32.5	10	25	
41-50	10	25	6	15	
Over 50	3	7.5	6	15	
<b>Sex</b>					
Men	18	45	15	37.5	Chi-squared test 0.462
Women	22	55	25	62.5	
<b>Marital status</b>					
Single	10	25	12	30	Fisher's test 0.726
Married	26	65	26	65	
Divorced	4	10	2	5	
<b>Occupational status</b>					
Employed by the government	4	10	2	5	Fisher's test 0.497
Construction worker	4	10	9	22.5	
Farmer	5	12.5	3	7.5	
Self-employed	16	40	13	32.5	
Housewife	11	27.5	13	32.5	
<b>Spinal disorders</b>					
Discopathy	38	95	39	97.5	Fisher's test 1
Abnormality	2	5	1	2.5	

**Table 2. The distribution of anxiety in the control and intervention groups during MRI**

Variable	Control		Intervention		Independent T-test
	Mean	Standard deviation	Mean	Standard deviation	
Anxiety experienced	37.4	15.41	23.32	12.72	P=0.000

reduced the MRI anxiety through applying relaxation and instrumental music, voiceless images, and telephone conversations [24]. Although the abovementioned methods are included as effective non-medicinal therapies the findings of which are consistent with those of the present study, their implementation is both costly and problematic. However, the essence of *Citrus aurantium* is both cheaper and easier to conduct than the aforementioned methods.

The findings of the present study are not consistent with those of the studies conducted by Holm et al., (Holm and Fitzmaurice [25]), Graham et al. [26], and Muzzarelli et al. [27]. In the study conducted by Holm et al, the inhalation of orange essence had no effect on the anxiety of children

hospitalized in the emergency department [25]. It seems that the lack of effectiveness of orange essence was owing to its inhalation method; they applied an electrical instrument for conducting aromatherapy. As it has been indicated by the researchers, most of the individuals in the aromatherapy group did not receive the essential oil. It is likely that that their method required a different distributor or more essence. Moreover, the hospital had a ventilation system that reduced the power of the essential oil. In the present study, a gauze attached to the patient's collar was used to avoid this problem. The findings of the study conducted by Graham et al, in Australia indicated that inhaling aromatherapy not only had no effect on reducing the anxiety of cancer patients undergoing radiotherapy but also

the anxiety of the control group was much less than that of the aromatherapy group [26]. Researchers maintain that the increased anxiety experienced by the aromatherapy patients is resulted from the relationship between the essential oil and the stressful experience of radiotherapy. In the study conducted by Muzzarelli et al. [27] aromatherapy (with a combination of lavender essential oil and grape seed oil) was reported to have no effect on reducing the patients' anxiety before conducting endoscopy [27]. This is not consistent with the findings of the present study. This inconsistency is possibly related to the amount of essential oil used in their study. The short inhalation time (5 minutes) and the concentration of the essential oil used (lavender 10% solved in grape seed oil) are possibly the causes for the insignificant results obtained. It is also likely that the difference between these studies and the present study is the nature of the diseases involved or the kind of inhalation essence. In terms of the exposure duration of the essence and its concentration, different studies have adopted different approaches. In their study, Namazi et al. [16], uses aromatherapy for reducing the anxiety of the first stage of the labor. In their study, they conducted aromatherapy with the *Citrus aurantium* essence 8% in 4 cc. This was repeated every 30 minutes [16]. In the present study, *Citrus aurantium* essence 10% was applied. It was used 10 minutes before scanning and the intervention was repeated during the scanning process. The percentage used and its duration calls for further studies.

The limitations of the present study were the existence of some environmental and psychological factors affecting the patient's anxiety; they were all out of the researcher's control. Moreover, it is likely that previous experience of being exposed to *Citrus aurantium* affect the findings of the present study as well. The main strong point of the present study was applying a specific tool of anxiety designed for MRI anxiety (Resonance Imaging-Anxiety Questionnaire (MRI-AQ)).

## 5. CONCLUSION

Given the findings of the present study, it was determined that using aromatherapy with *Citrus aurantium* is easier, inexpensive, and non-aggressive intervention, also likely to be effective in reducing MRI anxiety. Since the present study was conducted on patients during the MRI process, it is recommended to investigate the

effect of aromatherapy with *Citrus aurantium* during endoscopy and other diagnostic methods in future studies.

## ETHICAL APPROVAL AND CONSENT

The intended ethical considerations of the present study are explaining the purpose and nature of the study for all samples, acquiring informed consent from the samples, assuring the patients about the confidentiality of the data.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Harris LM, Menzies RG, Robinson J. Predictors of panic symptoms during magnetic resonance imaging scans. *International Journal of Behavioral Medicine*. 2001;8:80-87.
2. Kompel A, Murakami A, Guermazi A. Magnetic resonance imaging of nontraumatic musculoskeletal emergencies. *Magnetic Resonance Imaging Clinics*. 2016;24:369-389.
3. Thu H, Stutzman S, Supnet, Olson D. Factors associated with increased anxiety in the MRI waiting room. *Journal of Radiology Nursing*. 2015;34:170-4.
4. Chapman HA, Bernier D, Rusak B. MRI-related anxiety levels change within and between repeated scanning sessions. *Psychiatry Research: Neuroimaging*. 2010; 182:160-164.
5. Munn Z, Moola S, Lisy K, Riitano D, Murphy F. Claustrophobia in magnetic resonance imaging: A systematic review and meta-analysis. *Radiography*. 2015;21: e59-e63.
6. Dewey M, Schink T, Dewey CF. Claustrophobia during magnetic resonance imaging: Cohort study in over 55,000 patients. *Journal of Magnetic Resonance Imaging: An Official Journal of the International Society for Magnetic Resonance in Medicine*. 2007;26:1322-1327.
7. Enders J, Zimmermann E, Rief M, Martus P, Klingebiel R, Asbach P, Klessen C, Diederichs G, Bengner T, Teichgräber U. Reduction of claustrophobia during

- magnetic resonance imaging: methods and design of the "CLAUSTRO" randomized controlled trial. BMC Medical Imaging. 2011;11:4-15.
8. Kamrani F, Nazari M, Sahebalzamani M, Amin G. Effect of aromatherapy with lemon essential oil on anxiety after orthopedic surgery. Iranian Journal of Rehabilitation Research in Nursing. 2016;2:75-78.
  9. Tazegul G, Etcioğlu E, Yildiz F, Yildiz R, Tuney D. Can MRI related patient anxiety be prevented? Magnetic Resonance Imaging. 2015;33:180-183.
  10. Mohammed E, Atef J, Ellife H. Effectiveness of health instructions on reducing anxiety levels and claustrophobia among female adolescents undergoing Magnetic Resonance Imaging. American Journal of Research Communication. 2013;1:43-64.
  11. Tahmasbi H, Mahmoodi G, Mokhberi V, Hassani S, Akbarzadeh H, Rahnamai N. The impact of aromatherapy on the anxiety of patients experiencing coronary angiography. Zahedan J Res Med Sci. 2012;14:51-5.
  12. Sharifipour F, Mirmohammad Ali M, Hashemzadeh M. Comparison of the effect of *Citrus arantium* and *Salvia officinalis* aroma on post-cesarean section pain. The Iranian Journal of Obstetrics, Gynecology and Infertility. 2017;20:41-49.
  13. Najafi B, Mojab F, Ghaderi L, Farhadifar F, Roshani D, Seidi J. The effect of chamomile flower essence on pain severity after elective caesarean section under spinal anaesthesia: A randomized clinical trial. Journal of Clinical & Diagnostic Research. 2017;11:1-4.
  14. Najafi Z, Tagharrobi Z, Taghadosi M, Sharifi K, Farrokhian A. The effect of simultaneous aromatherapy and Quran recitation on anxiety level of patients with myocardial infarction. Complement Med J. 2014;4:700-12.
  15. Mojab F, Talebi A, Alavi Majd H. Investigating the effect of citrus aurantium aroma on severity of labor pain during active phase of labor. Complementary Medicine Journal of faculty of Nursing & Midwifery. 2014;4:733-744.
  16. Namazi M, Akbari SAA, Mojab F, Talebi A, Majd HA, Jannesari S. Effects of citrus aurantium (bitter orange) on the severity of first-stage labor pain. Iranian Journal of Pharmaceutical Research: IJPR. 2014;13: 10-11.
  17. Leite MP, Fassin JRJ, Baziloni EM, Almeida RN, Mattei R, Leite JR. Behavioral effects of essential oil of *Citrus aurantium* L. inhalation in rats. Revista Brasileira de Farmacognosia. 2008;18:661-666.
  18. Spielberger CD. STAI manual for the state-trait anxiety inventory. Self-Evaluation Questionnaire. 1970;1-24.
  19. Good M, Stiller C, Zauszniewski JA, Anderson GC, Stanton-Hicks M, Grass JA. Sensation and distress of pain scales: Reliability, validity, and sensitivity. Journal of Nursing Measurement. 2001;9:219-238.
  20. Abrosh Z, Rezai M, Ashrafi F. Effect of antibacterial activity of *Salvia officinalis* essence. J Med Aromatic Plants. 2004;20: 457-68.
  21. Ahlander BM, Aarestedt K, Engvall J, Maret E, Ericsson E. Development and validation of a questionnaire evaluating patient anxiety during Magnetic Resonance imaging: The magnetic resonance imaging-anxiety questionnaire (MRI-AQ). Journal of Advanced Nursing. 2016;72:1368-1380.
  22. Sharifipour F, Baigi SS, Mirmohammadali M. The aromatic effect of *Citrus arantium* on pain and vital signs after cesarean section. International Journal of Biology, Pharmacy, and Allied Sciences. 2015;4: 5063-5072.
  23. Akhlaghi M, Shabaniyan G, Rafieian-Kopaei M, Parvin N, Saadat M, Akhlaghi M. *Citrus aurantium* blossom and preoperative anxiety. Revista Brasileira de Anestesiologia. 2011;61:707-712.
  24. Tugwell JR, Goulden N, Mullins P. Alleviating anxiety in patients prior to MRI: A pilot single-centre single-blinded randomised controlled trial to compare video demonstration or telephone conversation with a radiographer versus routine intervention. Radiography. 2018; 24:122-129.
  25. Holm L, Fitzmaurice L. Emergency department waiting room stress: Can music or aromatherapy improve anxiety scores? Pediatric Emergency Care. 2008; 24:836-838.
  26. Graham P, Browne L, Cox H, Graham J. Inhalation aromatherapy during radiotherapy: Results of a placebo-

- controlled double-blind randomized trial. Journal of Clinical Oncology. 2003;21: 2372-2376.
27. Muzzarelli L, Force M, Sebold M. Aromatherapy and reducing preprocedural anxiety: A controlled prospective study. Gastroenterology Nursing: The Official Journal of the Society of Gastroenterology Nurses and Associates. 2006;29:466-471.

---

© 2019 Ahani et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/46877>